

Characterization of the Anti-Bovine Podoplanin Monoclonal Antibody PMab-44

Shinji Yamada,¹ Ryusuke Honma,² Mika K. Kaneko,¹ Takuro Nakamura,¹ Miyuki Yanaka,¹ Noriko Saidoh,¹ Michiaki Takagi,² Satoru Konnai,³ and Yukinari Kato^{1,4}

A type I transmembrane sialoglycoprotein podoplanin (PDPN) is expressed in several normal cells, including podocytes of the kidney, type I alveolar cells of the lung, and lymphatic endothelial cells. We recently produced an anti-bovine PDPN (bovPDPN) monoclonal antibody (mAb), PMab-44, by immunizing mice with recombinant proteins of bovPDPN. In this study, we determined the critical epitope of PMab-44 for the recognition of bovPDPN using many deletion mutants and point mutants of bovPDPN. Flow cytometric analyses revealed that the epitope of PMab-44 was Glu46-Thr50, which corresponds to platelet aggregation-stimulating (PLAG) domain-3. The important amino acids in the PMab-44 epitope were determined to be Glu46, Tyr48, and Thr50. Western blot analysis also confirmed these results, indicating that the PLAG domain of bovPDPN is also important in immunogenicity for producing useful anti-PDPN mAbs.

Keywords: podoplanin, bovine PDPN, PMab-44

Introduction

A TYPE I TRANSMEMBRANE sialoglycoprotein podoplanin (PDPN), also known as Aggrus,⁽¹⁾ activates platelet aggregation by binding to the C-type lectin-like receptor-2 (CLEC-2) on platelets.⁽²⁻⁴⁾ The interaction between PDPN and CLEC-2 was mainly observed at Glu47 and Asp48 in the platelet aggregation-stimulating (PLAG) domain and the α 2,6-linked sialic acid at Thr52 of human PDPN.^(2,4,5) This interaction facilitates blood/lymphatic vessel separation.⁽⁶⁾ The actin cytoskeleton is rearranged in CLEC-2-expressing dendritic cells to promote efficient motility along PDPN-expressing stromal surfaces.⁽⁷⁾ The signal from CLEC-2 to PDPN controls the contractility of fibroblastic reticular cells and lymph node microarchitecture.⁽⁸⁾ The physical elasticity of lymph nodes is maintained by the PDPN of stromal fibroblastic reticular cells.⁽⁹⁾

PDPN is expressed in several normal cells, including lymphatic endothelial cells, renal podocytes, pulmonary type I alveolar cells, osteocytes, and chondrocytes.⁽¹⁰⁾ The expression of PDPN has been reported in many malignant tumors, such as oral squamous cell carcinomas,⁽¹¹⁾ malignant brain tumors,⁽¹²⁻¹⁵⁾ lung cancers,⁽¹⁶⁾ esophageal cancers,⁽¹⁷⁾ malignant mesotheliomas,^(18,19) testicular tumors,⁽²⁰⁾ osteosarcomas,^(10,21,22) and chondrosarcomas.⁽¹⁰⁾ PDPN expres-

sion is also associated with malignant progression and cancer metastasis.^(2,12,23)

We had previously established the anti-bovine PDPN (bovPDPN) monoclonal antibody (mAb) PMab-44 (IgG₁, kappa), which was useful for detecting podocytes and lymphatic endothelial cells in normal bovine tissues in immunohistochemistry.⁽²⁴⁾ PMab-44 also detected bovPDPN specifically in flow cytometry. Our previous research revealed that bovPDPN lacks platelet aggregation-inducing activity because the predicted amino acid sequence of bovPDPN has a sporadic deletion in the PLAG domain.⁽²⁵⁾ Therefore, the functional character of bovPDPN may be somewhat different from other mammalian PDPN. However, bovPDPN has so far attracted little attention apart from a few studies.^(26,27)

In this study, we further determined the epitope of PMab-44 in flow cytometry and western blotting. This information could be useful in bovPDPN veterinary research.

Materials and Methods

Cell lines

Chinese hamster ovary (CHO)-K1 was purchased from the American Type Culture Collection (ATCC, Manassas, VA). CHO-K1 cells were transfected with the bovPDPN-MAP tag⁽²⁸⁾ plasmid using a Gene Pulser Xcell electroporation

¹Department of Antibody Drug Development, Tohoku University Graduate School of Medicine, Sendai, Japan.

²Department of Orthopedic Surgery, Yamagata University Faculty of Medicine, Yamagata, Japan.

³Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo, Japan.

⁴New Industry Creation Hatchery Center, Tohoku University, Sendai, Japan.

system (Bio-Rad Laboratories, Inc., Berkeley, CA). CHO-K1 and CHO/bovPDPN were cultured in RPMI 1640 medium (Nacalai Tesque, Inc., Kyoto, Japan), supplemented with 10% heat-inactivated fetal bovine serum (Thermo Fisher Scientific, Inc., Waltham, MA), 100 units/mL of penicillin, 100 μ g/mL of streptomycin, and 25 μ g/mL of amphotericin B (Nacalai Tesque, Inc.) at 37°C in a humidified atmosphere of 5% CO₂ and 95% air.

Production of bovPDPN mutants

The amplified bovPDPN cDNA was subcloned into a pCAG vector (Wako Pure Chemical Industries Ltd., Osaka, Japan) and a MAP tag⁽²⁸⁾ was added at the N-terminus. Deletion mutation of a bovPDPN sequence was performed using a HotStar HiFidelity polymerase chain reaction (PCR; Qiagen, Inc., Hilden, Germany) with oligonucleotides containing the desired mutations. PCR fragments bearing the desired mutations were inserted into a pCAG vector using the In-Fusion PCR Cloning Kit (Clontech, Palo Alto, CA). Substitutions of amino acids to alanine in bovPDPN sequences were conducted using the QuikChange lightning site-directed mutagenesis kit (Agilent Technologies, Inc., Santa Clara, CA). CHO-K1 cells were transfected with the plasmids using a Gene Pulser Xcell electroporation system (Bio-Rad Laboratories, Inc.).

Flow cytometry

Cells were harvested by brief exposure to 0.25% trypsin/1 mM EDTA (Nacalai Tesque, Inc.). After washing with 0.1% BSA/PBS, the cells were treated with PMab-44 (10 μ g/mL) and anti-MAP tag⁽²⁸⁾ (clone: PMab-1, 1 μ g/mL) for 30 minutes at 4°C followed by treatment with Oregon green-conjugated anti-mouse IgG or anti-rat IgG (1:1000 diluted; Thermo Fisher Scientific, Inc.). Fluorescence data were collected using a Cell Analyzer EC800 (Sony Corp., Tokyo, Japan). PMab-1 was previously developed against mouse PDPN⁽²⁹⁾ and has been utilized as an anti-MAP tag mAb.⁽²⁸⁾ PMab-1 does not crossreact with other PDPN proteins such as bovPDPN,⁽²⁴⁾ dog PDPN,^(30–32) rat PDPN,⁽³³⁾ rabbit PDPN,^(34,35) and human PDPN.⁽²⁹⁾

Western blot analyses

Cell lysates (10 μ g) were boiled in sodium dodecyl sulfate (SDS) sample buffer (Nacalai Tesque, Inc.). The proteins were electrophoresed on 5%–20% polyacrylamide gels (Wako Pure Chemical Industries Ltd.) and were transferred onto a PVDF membrane (Merck KGaA, Darmstadt, Germany). After blocking with 4% skim milk (Nacalai Tesque, Inc.), the membrane was incubated with PMab-44, anti-MAP tag⁽²⁸⁾ (clone: PMab-1), and anti- β -actin (clone AC-15; Sigma-Aldrich Corp., St. Louis, MO) and then with





FIG. 1. Epitope mapping of an anti-bovPDPN mAb PMab-44 using deletion mutants by flow cytometry. **(A)** Deletion mutants of bovPDPN (dN23, dN40, dN60, dN80, and dN100) were treated with PMab-44 (10 μ g/mL) and PMab-1 (1 μ g/mL) for 30 minutes at 4°C followed by treatment with Oregon green-conjugated anti-mouse IgG or anti-rat IgG. **(B)** Deletion mutants of bovPDPN (dN23, dN45, dN50, and dN55) were treated with PMab-44 (10 μ g/mL) and PMab-1 (1 μ g/mL) for 30 minutes at 4°C followed by treatment with Oregon green-conjugated anti-mouse IgG or anti-rat IgG. Fluorescence data were collected using a Cell Analyzer EC800. Red line: PMab-44 or anti-PA tag. Black line: negative control. bovPDPN, bovine PDPN; mAb, monoclonal antibody; PDPN, podoplanin.



peroxidase-conjugated anti-mouse or anti-rat antibodies (1:1000 diluted; Dako; Agilent Technologies, Inc.). Subsequently, the membrane was developed with the Pierce Western Blotting Substrate Plus (Thermo Fisher Scientific, Inc.) or the ImmunoStar LD Chemiluminescence Reagent (Wako Pure Chemical Industries Ltd.) using a Sayaca-Imager (DRC Co. Ltd., Tokyo, Japan).

Determination of the binding affinity using flow cytometry

CHO/bovPDPN (2×10^5 cells) were resuspended with 100 μ L of serially diluted antibodies (0.6 ng/mL–10 μ g/mL) followed by secondary anti-mouse IgG (Thermo Fisher Scientific, Inc.). Fluorescence data were collected using a cell analyzer (EC800; Sony Corp.). The dissociation constants (K_D) were obtained by fitting the binding isotherms using the built-in one-site binding models in GraphPad PRISM 6 (GraphPad Software, Inc., La Jolla, CA).

Results and Discussion

Epitope mapping of an anti-bovPDPN mAb PMab-44 by flow cytometry

We previously showed the usefulness of an anti-bovPDPN mAb PMab-44 in immunohistochemical analysis⁽²⁴⁾; in this study we performed the epitope mapping using flow cytometry and western blot analysis.

First, we produced five N-terminal deletion mutants of bovPDPN: dN23 (23–169 amino acids), dN40 (40–169 amino acids), dN60 (60–169 amino acids), dN80 (80–169 amino acids), and dN100 (100–169 amino acids). Flow cytometric analysis revealed that PMab-44 reacted with dN23 and dN40; in contrast, it did not react with dN60, dN80, and dN100 (Fig. 1A). The anti-MAP tag PMab-1 recognized all deletion mutants. These results indicate that the N-terminal of the PMab-44 epitope exists between Lys40 and Ala60 of bovPDPN.⁽²⁵⁾ Furthermore, we produced three N-terminal deletion mutants of bovPDPN: dN45 (45–169 amino acids), dN50 (50–169 amino acids), and dN55 (55–169 amino acids). Flow cytometric analysis revealed that PMab-44 reacted with dN23 and dN45; in contrast, it did not react with dN50 and dN55 (Fig. 1B). These results indicate that the N-terminal of the PMab-44 epitope exists between Val45 and Thr50 of bovPDPN.⁽²⁵⁾

Next, we produced 10 point mutants of bovPDPN: V45A, E46A, D47A, Y48A, T49A, T50A, T51A, P52A, A53G, and A54G. Figure 2 shows that two point mutants, such as E46A and Y48A, completely lost the reaction by PMab-44, and T50A showed little reaction by PMab-44. All point mutants were recognized by the anti-MAP tag PMab-1. These results

FIG. 2. Epitope mapping of an anti-bovPDPN mAb PMab-44 using point mutants by flow cytometry. Point mutants of bovPDPN (V45A, E46A, D47A, Y48A, T49A, T50A, T51A, P52A, A53G, and A54G) were treated with PMab-44 (10 μ g/mL) and anti-MAP tag (1 μ g/mL) for 30 minutes at 4°C followed by treatment with Oregon green-conjugated anti-mouse IgG or anti-rat IgG. Fluorescence data were collected using a Cell Analyzer EC800. Red line: PMab-44 or anti-PA tag. Black line: negative control.

indicate that three amino acids of bovPDPN, Glu46, Tyr48, and Thr50, are critical for PMAb-44 recognition.

Epitope mapping of an anti-bovPDPN mAb PMAb-44 by western blot analysis

We next performed western blot analysis to confirm the results of epitope mapping of PMAb-44 by flow cytometry. As shown in Figure 3A, PMAb-44 did not react with V45A, E46A, D47A, Y48A, T49A, and T50A of bovPDPN. In contrast, PMAb-44 reacted with T51A, P52A, A53G, and A54G. All point mutants were detected by the anti-MAP tag PMAb-1. These results indicated that Val45, Glu46, Asp47, Tyr48, Thr49, and Thr50 are important in the recognition of PMAb-44 against bovPDPN in western blot analysis. The epitope of PMAb-44 is illustrated in Figure 3B. The sequence of the PMAb-44 epitope (VEDYTT) shows low homology with the corresponding peptide sequence of human PDPN (AEDDVV), indicating that PMAb-44 does not react with human PDPN.⁽²⁴⁾

Determination of the binding affinity using flow cytometry

We next performed a kinetic analysis of the interaction of PMAb-44 with CHO/bovPDPN using flow cytometry. As




FIG. 3. Epitope mapping of an anti-bovPDPN mAb PMAb-44 using point mutants by western blot analysis. **(A)** Cell lysate (10 μ g) was electrophoresed on 5%–20% polyacrylamide gels and transferred onto a PVDF membrane. After blocking, the membrane was incubated with 1 μ g/mL of mAbs (PMAb-44, anti-MAP tag, and anti- β -actin) and then with peroxidase-conjugated secondary antibodies; the membrane was developed with Pierce Western Blotting Substrate Plus or ImmunoStar LD Chemiluminescence Reagent and detected using a Sayaca-Imager. **(B)** Schematic illustration of the epitope of PMAb-44. Gray shade: identical amino acid residue between bovPDPN and human PDPN. Arrow head: conserved Thr of PLAG domain. Red character: PMAb-44 epitope. PLAG domain, platelet aggregation-stimulating domain.




FIG. 4. Determination of binding affinity using flow cytometry. CHO/bovPDPN cells were resuspended with 100 μ L of serially diluted antibodies (0.6 ng/mL–10 μ g/mL) followed by secondary anti-mouse IgG. Fluorescence data were collected using a cell analyzer. CHO, Chinese hamster ovary.

shown in Figure 4, the K_D of PMAb-44 was determined as 3.3×10^{-9} M, indicating that PMAb-44 possesses a high affinity against bovPDPN.

In conclusion, the most important amino acids for PMAb-44 recognition against bovPDPN were determined to be Glu46, Tyr48, and Thr50; second in importance were Val45, Asp47, and Thr49. Three amino acids, Glu46, Tyr48, and Thr50, exist in the PLAG3 of bovPDPN,⁽²⁵⁾ indicating that the PLAG3 of bovPDPN is also important for immunogenicity in producing useful anti-PDPN mAbs.

Acknowledgments

This research was supported in part by grants from the Science and Technology Research Promotion Program for Agriculture, Forestry, Fisheries, and Food Industry, Japan (grant 26058B to S.K.), and the NARO, Bio-oriented Technology Research Advancement Institution (the special scheme project on regional developing strategy: grant 16817557 to S.K.). This work was also supported in part by Project for utilizing glycans in the development of innovative drug discovery technologies from Japan Agency for Medical Research and Development, AMED (Y.K.), by the Basic Science and Platform Technology Program for Innovative Biological Medicine from AMED (Y.K.), by the Platform for Drug Discovery, Informatics, and Structural Life Science (PDIS) from AMED (Y.K.), by the Regional Innovation Strategy Support Program from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan (Y.K.), by JSPS KAKENHI Grant Number 26440019 and 17K07299 (M.K.K.) and 16K10748 (Y.K.). This work was performed in part under the Cooperative Research Program of Institute for Protein Research, Osaka University, CR-16-05 and CR-17-05 and by the Grant for Joint Research Project of the Institute of Medical Science, the University of Tokyo. The authors would like to thank Enago (www.enago.jp) for the English language review.

Author Disclosure Statement

No competing financial interests exist.

References

- Kato Y, Fujita N, Kunita A, Sato S, Kaneko M, Osawa M, and Tsuruo T: Molecular identification of Aggrus/T1alpha as a platelet aggregation-inducing factor expressed in colorectal tumors. *J Biol Chem* 2003;278:51599–51605.
- Kato Y, Kaneko MK, Kunita A, Ito H, Kameyama A, Ogasawara S, Matsuura N, Hasegawa Y, Suzuki-Inoue K, Inoue O, Ozaki Y, and Narimatsu H: Molecular analysis of the pathophysiological binding of the platelet aggregation-inducing factor podoplanin to the C-type lectin-like receptor CLEC-2. *Cancer Sci* 2008;99:54–61.
- Kaneko MK, Kunita A, Abe S, Tsujimoto Y, Fukayama M, Goto K, Sawa Y, Nishioka Y, and Kato Y: Chimeric anti-podoplanin antibody suppresses tumor metastasis through neutralization and antibody-dependent cellular cytotoxicity. *Cancer Sci* 2012;103:1913–1919.
- Suzuki-Inoue K, Kato Y, Inoue O, Kaneko MK, Mishima K, Yatomi Y, Yamazaki Y, Narimatsu H, and Ozaki Y: Involvement of the snake toxin receptor CLEC-2, in podoplanin-mediated platelet activation, by cancer cells. *J Biol Chem* 2007;282:25993–26001.
- Nagae M, Morita-Matsumoto K, Kato M, Kaneko MK, Kato Y, and Yamaguchi Y: A platform of C-type lectin-like receptor CLEC-2 for binding O-glycosylated podoplanin and nonglycosylated rhodocytin. *Structure* 2014;22:1711–1721.
- Bertozzi CC, Schmaier AA, Mericko P, Hess PR, Zou Z, Chen M, Chen CY, Xu B, Lu MM, Zhou D, Sebzda E, Santore MT, Merianos DJ, Stadtfeld M, Flake AW, Graf T, Skoda R, Maltzman JS, Koretzky GA, and Kahn ML: Platelets regulate lymphatic vascular development through CLEC-2-SLP-76 signaling. *Blood* 2010;116:661–670.
- Acton SE, Astarita JL, Malhotra D, Lukacs-Kornek V, Franz B, Hess PR, Jakus Z, Kuligowski M, Fletcher AL, Elpek KG, Bellemare-Pelletier A, Sceats L, Reynolds ED, Gonzalez SF, Graham DB, Chang J, Peters A, Woodruff M, Kim YA, Swat W, Morita T, Kuchroo V, Carroll MC, Kahn ML, Wucherpfennig KW, and Turley SJ: Podoplanin-rich stromal networks induce dendritic cell motility via activation of the C-type lectin receptor CLEC-2. *Immunity* 2012;37:276–289.
- Astarita JL, Cremasco V, Fu J, Darnell MC, Peck JR, Nieves-Bonilla JM, Song K, Kondo Y, Woodruff MC, Gogineni A, Onder L, Ludewig B, Weimer RM, Carroll MC, Mooney DJ, Xia L, and Turley SJ: The CLEC-2-podoplanin axis controls the contractility of fibroblastic reticular cells and lymph node microarchitecture. *Nat Immunol* 2015;16:75–84.
- Acton SE, Farrugia AJ, Astarita JL, Mourao-Sa D, Jenkins RP, Nye E, Hooper S, van Blijswijk J, Rogers NC, Snelgrove KJ, Rosewell I, Moita LF, Stamp G, Turley SJ, Sahai E, and Reis e Sousa C: Dendritic cells control fibroblastic reticular network tension and lymph node expansion. *Nature* 2014;514:498–502.
- Ariizumi T, Ogose A, Kawashima H, Hotta T, Li G, Xu Y, Umezumi H, Sugai M, and Endo N: Expression of podoplanin in human bone and bone tumors: New marker of osteogenic and chondrogenic bone tumors. *Pathol Int* 2010;60:193–202.
- Ochoa-Alvarez JA, Krishnan H, Pastorino JG, Nevel E, Kephart D, Lee JJ, Retzbach EP, Shen Y, Fatahzadeh M, Baredes S, Kalyoussef E, Honma M, Adelson ME, Kaneko MK, Kato Y, Young MA, Deluca-Rapone L, Shienbaum AJ, Yin K, Jensen LD, and Goldberg GS: Antibody and lectin target podoplanin to inhibit oral squamous carcinoma cell migration and viability by distinct mechanisms. *Oncotarget* 2015;6:9045–9060.
- Mishima K, Kato Y, Kaneko MK, Nishikawa R, Hirose T, and Matsutani M: Increased expression of podoplanin in malignant astrocytic tumors as a novel molecular marker of malignant progression. *Acta Neuropathol (Berl)* 2006;111:483–488.
- Mishima K, Kato Y, Kaneko MK, Nakazawa Y, Kunita A, Fujita N, Tsuruo T, Nishikawa R, Hirose T, and Matsutani M: Podoplanin expression in primary central nervous system germ cell tumors: A useful histological marker for the diagnosis of germinoma. *Acta Neuropathol (Berl)* 2006;111:563–568.
- Kato Y, Vaidyanathan G, Kaneko MK, Mishima K, Srivastava N, Chandramohan V, Pegram C, Keir ST, Kuan CT, Bigner DD, and Zalutsky MR: Evaluation of anti-podoplanin rat monoclonal antibody NZ-1 for targeting malignant gliomas. *Nucl Med Biol* 2010;37:785–794.
- Kato Y, and Kaneko MK: A cancer-specific monoclonal antibody recognizes the aberrantly glycosylated podoplanin. *Sci Rep* 2014;4:5924.
- Kato Y, Kaneko M, Sata M, Fujita N, Tsuruo T, and Osawa M: Enhanced expression of Aggrus (T1alpha/podoplanin), a platelet-aggregation-inducing factor in lung squamous cell carcinoma. *Tumor Biol* 2005;26:195–200.
- Schoppmann SF, Jesch B, Riegler MF, Maroske F, Schwameis K, Jomrich G, and Birner P: Podoplanin expressing cancer associated fibroblasts are associated with unfavourable prognosis in adenocarcinoma of the esophagus. *Clin Exp Metastasis* 2013;30:441–446.
- Kimura N, and Kimura I: Podoplanin as a marker for mesothelioma. *Pathol Int* 2005;55:83–86.
- Abe S, Morita Y, Kaneko MK, Hanibuchi M, Tsujimoto Y, Goto H, Kakiuchi S, Aono Y, Huang J, Sato S, Kishuku Y, Taniguchi Y, Azuma M, Kawazoe K, Sekido Y, Yano S, Akiyama S, Sone S, Minakuchi K, Kato Y, and Nishioka Y: A novel targeting therapy of malignant mesothelioma using anti-podoplanin antibody. *J Immunol* 2013;190:6239–6249.
- Kato Y, Sasagawa I, Kaneko M, Osawa M, Fujita N, and Tsuruo T: Aggrus: A diagnostic marker that distinguishes seminoma from embryonal carcinoma in testicular germ cell tumors. *Oncogene* 2004;23:8552–8556.
- Kaneko MK, Oki H, Ogasawara S, Takagi M, and Kato Y: Anti-podoplanin monoclonal antibody LpMab-7 detects metastatic lesions of osteosarcoma. *Monoclon Antib Immunodiagn Immunother* 2015;34:154–161.
- Kunita A, Kashima TG, Ohazama A, Grigoriadis AE, and Fukayama M: Podoplanin is regulated by AP-1 and promotes platelet aggregation and cell migration in osteosarcoma. *Am J Pathol* 2011;179:1041–1049.
- Kunita A, Kashima TG, Morishita Y, Fukayama M, Kato Y, Tsuruo T, and Fujita N: The platelet aggregation-inducing factor aggrus/podoplanin promotes pulmonary metastasis. *Am J Pathol* 2007;170:1337–1347.
- Honma R, Ogasawara S, Kaneko M, Fujii Y, Oki H, Nakamura T, Takagi M, Konnai S, and Kato Y: PMab-44 Detects Bovine Podoplanin in Immunohistochemistry. *Monoclon Antib Immunodiagn Immunother* 2016;35:186–190.
- Kaneko MK, Kato Y, Kitano T, and Osawa M: Conservation of a platelet activating domain of Aggrus/podoplanin

- as a platelet aggregation-inducing factor. *Gene* 2006;378: 52–57.
26. Gao Y, Qin L, Yang Y, Dong X, Zhao Z, Zhang G, and Zhao Z: PDPN gene promotes the proliferation of immature Bovine Sertoli cells in vitro. *Anim Reprod Sci* 2017;179: 35–43.
 27. Nitta A, Shirasuna K, Nibuno S, Bollwein H, Shimizu T, and Miyamoto A: Downregulation of lymphatic vessel formation factors in PGF2alpha-induced luteolysis in the cow. *J Reprod Dev* 2013;59:296–301.
 28. Fujii Y, Kaneko MK, and Kato Y: MAP Tag: A Novel Tagging System for Protein Purification and Detection. *Monoclon Antib Immunodiagn Immunother* 2016;35:293–299.
 29. Kaji C, Tsujimoto Y, Kato Kaneko M, Kato Y, and Sawa Y: Immunohistochemical examination of novel rat monoclonal antibodies against mouse and human podoplanin. *Acta Histochem Cytochem* 2012;45:227–237.
 30. Honma R, Kaneko MK, Ogasawara S, Fujii Y, Konnai S, Takagi M, and Kato Y: Specific detection of dog podoplanin expressed in renal glomerulus by a novel monoclonal antibody PMab-38 in immunohistochemistry. *Monoclon Antib Immunodiagn Immunother* 2016;35:212–216.
 31. Kaneko MK, Honma R, Ogasawara S, Fujii Y, Nakamura T, Saidoh N, Takagi M, Kagawa Y, Konnai S, and Kato Y: PMab-38 recognizes canine podoplanin of squamous cell carcinomas. *Monoclon Antib Immunodiagn Immunother* 2016;35:263–266.
 32. Ogasawara S, Honma R, Kaneko MK, Fujii Y, Kagawa Y, Konnai S, and Kato Y: Podoplanin expression in canine melanoma. *Monoclon Antib Immunodiagn Immunother* 2016;35:304–306.
 33. Oki H, Honma R, Ogasawara S, Fujii Y, Liu X, Takagi M, Kaneko MK, and Kato Y: Development of sensitive monoclonal antibody PMab-2 against rat podoplanin. *Monoclon Antib Immunodiagn Immunother* 2015;34:396–403.
 34. Honma R, Fujii Y, Ogasawara S, Oki H, Liu X, Nakamura T, Kaneko MK, Takagi M, and Kato Y: Establishment of a novel monoclonal antibody PMab-32 against rabbit podoplanin. *Monoclon Antib Immunodiagn Immunother* 2016; 35:41–47.
 35. Honma R, Fujii Y, Ogasawara S, Oki H, Konnai S, Kagawa Y, Takagi M, Kaneko MK, and Kato Y: Critical epitope of anti-rabbit podoplanin monoclonal antibodies for immunohistochemical analysis monoclon. *Antib Immunodiagn Immunother* 2016;35:65–72.

Address correspondence to:

*Dr. Yukinari Kato
New Industry Creation Hatchery Center
Tohoku University
Department of Antibody Drug Development
Tohoku University Graduate School of Medicine
2-1 Seiryomachi, Aoba-ku
Sendai 980-8575
Japan*

*E-mail: yukinari-k@bea.hi-ho.ne.jp;
yukinarikato@med.tohoku.ac.jp*

Received: March 9, 2017

Accepted: April 16, 2017