

Open camera or QR reader and
scan code to access this article
and other resources online.



Establishment of a Novel Anti-Human CCR6 Monoclonal Antibody C₆Mab-19 with the High Binding Affinity in Flow Cytometry

Tomohiro Tanaka,* Kaishi Kitamura,* Hiroyuki Suzuki,* Mika K. Kaneko, and Yukinari Kato

CC chemokine receptor 6 (CCR6) is a member of the G-protein-coupled receptor family that is highly expressed in B lymphocytes, effector and memory T cells, regulatory T cells, and immature dendritic cells. CCR6 has been revealed to have important functions in many pathological conditions, such as cancer, intestinal bowel disease, psoriasis, and autoimmune diseases. The only CCR6 chemokine ligand, CC motif chemokine ligand 20 (CCL20), is also involved in pathogenesis by interacting with CCR6. The CCL20/CCR6 axis is drawing attention as an attractive therapeutic target for various diseases. In this study, we developed novel monoclonal antibodies (mAbs) against human CCR6 (hCCR6) using the peptide immunization method, which are applicable to flow cytometry and immunohistochemistry. The established anti-hCCR6 mAb, clone C₆Mab-19 (mouse IgG₁, kappa), reacted with hCCR6-overexpressed Chinese hamster ovary-K1 (CHO/hCCR6), human liver carcinoma (HepG2), and human differentiated hepatoma (HuH-7) cells in flow cytometry. The dissociation constant (K_D) of C₆Mab-19 was determined as 3.0×10^{-10} M for CHO/hCCR6, 6.9×10^{-10} M for HepG2, and 1.8×10^{-10} M for HuH-7. Thus, C₆Mab-19 could bind to exogenously and endogenously expressed hCCR6 with extremely high affinity. Furthermore, C₆Mab-19 could stain formalin-fixed paraffin-embedded lymph node tissues from a patient with non-Hodgkin lymphoma by immunohistochemistry. Therefore, C₆Mab-19 is suitable for detecting hCCR6-expressing cells and tissues and could be useful for pathological analysis and diagnosis.

Keywords: CCR6, monoclonal antibody, peptide immunization, flow cytometry, immunohistochemistry

Introduction

CHEMOKINES ARE CHEMOTACTIC cytokines with a small molecular weight (8,000–12,000) and are classified into four categories: CC, CXC, CX3C, and XC subfamilies, according to the number and position of cysteine residues in their N-terminus.^{1,2} Approximately 50 chemokines, with diverse functions, have been discovered so far. Chemokines were originally thought to function in the leukocyte development, differentiation, and migration, but they are now known to be deeply involved in the pathogenesis of numerous diseases, including inflammatory diseases and cancers.^{3,4} Chemokines exert their functions by binding to corresponding seven-transmembrane G-protein-coupled receptors

(GPCRs). This chemokine-GPCR interaction is important for the physiological functions and pathogenesis of cells.⁵

In 1997, the CC chemokine receptor 6 (CCR6) was discovered as a specific receptor for CC motif chemokine ligand 20 (CCL20).⁶ CCR6 has been reported to be expressed in tissues, including intestinal mucosa, lung mucosa, and lymph nodes, as well as B and T cells. CCL20, also known as macrophage inflammatory protein-3 α or liver and activation-regulated chemokine, is expressed in tissues, such as the liver, lung, and lymph nodes. CCL20 is expressed in immune-related cells, such as dendritic cells, Th17 cells, natural killer cells, and B cells.⁴ CCL20 functions in cooperation with CCR6.

The binding mode of CCL20 and CCR6 has been structurally elucidated.⁷ CCL20 has the shortest N-terminus

Department of Antibody Drug Development, Tohoku University Graduate School of Medicine, Sendai, Japan.

This article was previously published in preprint.org (doi: 10.20944/preprints202303.0181.v1).

*These authors contributed equally to this work.

among chemokines and binds to a shallow surface region of CCR6. CCL20 and/or CCR6 are involved in the progression of numerous pathologies.⁸ CCL20/CCR6-activated M1 macrophages cause adenoid hypertrophy by adenoid epithelium inflammation. An imiquimod-induced psoriasis model reported that skin-homing CCR6⁺ $\gamma\delta^{\text{Low}}$ T cells strongly produce interleukin-17A and cause inflammation.⁹ Furthermore, a CCR6 antagonist ameliorates inflammatory bowel disease by regulating immune cell chemotaxis.¹⁰ These reports indicate that CCL20/CCR6 contributes to pathogenesis by controlling immune cells.

Many reports have shown the effects of CCL20/CCR6 in tumors.¹¹ The regulatory T (Treg) cells, which act as a suppressive factor of cancer immunity, interact with CCL20 in the tumor microenvironment via CCR6 on Treg and contribute to cancer progression and poor prognoses, such as oral, head and neck squamous carcinomas, and hepatocellular carcinomas.^{12–14} CCL20, which is secreted by macrophages and stromal fibroblasts, promotes cancer cells or Th17 cell behavior.^{15–17} CCL20/CCR6 appears to play roles in managing immune cells around tumors. CCR6-expressing chimeric antigen receptor-T cells (CAR-T) have been shown to be effective in lung cancer models.¹⁸ These findings make the CCL20/CCR6 axis an attractive therapeutic target for various diseases.

Previously, we have developed various monoclonal antibodies (mAbs) against mouse and human chemokine receptors, including an anti-mouse CCR2 mAb,¹⁹ anti-mouse CCR3 mAbs,^{20–23} an anti-mouse CCR4 mAb,²⁴ anti-mouse CCR6 mAbs,²⁵ anti-mouse CCR8 mAbs,²⁶ an anti-mouse CCR9 mAb,²⁷ an anti-mouse CXCR6 mAb,²⁸ an anti-human CCR2 mAb,²⁹ and anti-human CCR9 mAbs.^{30,31} In this study, we developed anti-human CCR6 (hCCR6) mAbs using the N-terminal peptide immunization method.

Materials and Methods

Cell lines

Human liver carcinoma (HepG2) and human differentiated hepatoma (HuH-7) were obtained from the Cell Resource Center for Biomedical Research Institute of Development, Aging and Cancer at Tohoku University. Chinese hamster ovary (CHO)-K1 and P3X63Ag8U.1 (P3U1) were obtained from the American Type Culture Collection (Manassas, VA). The expression plasmid of hCCR6 (pCMV6neo-hCCR6-Myc-DDK; Accession No.: NM_004367) was purchased from OriGene Technologies, Inc. (Rockville, MD). A Neon Transfection System (Thermo Fisher Scientific, Inc., Waltham, MA) was used to transfect the hCCR6 plasmid into CHO-K1 cells. Stable transfectants were established through cell sorting using a cell sorter (SH800; Sony Corp., Tokyo, Japan) and cultivation in a medium containing 0.5 mg/mL of G418 (Nacalai Tesque, Inc., Kyoto, Japan).

CHO-K1, P3U1, hCCR6-overexpressed CHO-K1 (CHO/hCCR6), and HuH-7 were cultured in the Roswell Park Memorial Institute (RPMI)-1640 medium (Nacalai Tesque, Inc.). HepG2 cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Nacalai Tesque, Inc.). RPMI-1640 medium and DMEM were supplemented with 10% heat-inactivated fetal bovine serum (FBS; Thermo Fisher Scientific, Inc.), 100 units/mL penicillin, 100 $\mu\text{g}/\text{mL}$ streptomycin, and 0.25 $\mu\text{g}/\text{mL}$ amphotericin B (Nacalai

Tesque, Inc.). Cells were grown in a humidified incubator at 37°C with an atmosphere of 5% CO₂ and 95% air.

Antibodies

An anti-hCCR6 mAb (clone G034E3) was purchased from BioLegend (San Diego, CA). Secondary Alexa Fluor 488-conjugated anti-mouse IgG was purchased from Cell Signaling Technology, Inc. (Danvers, MA).

Peptides

A partial sequence of the N-terminal extracellular region of hCCR6 (1-MNFSDFVDSSEYDFVSVNT-19) plus C-terminal cysteine (hCCR6p1-19C) was synthesized by Eurofins Genomics KK (Tokyo, Japan). Subsequently, the keyhole limpet hemocyanin (KLH) was conjugated at the C-terminus of the peptide (hCCR6p1-19C-KLH).

Production of hybridomas

The two 5-week-old BALB/c mice were purchased from CLEA Japan (Tokyo, Japan) and were housed under specific pathogen-free conditions. All animal experiments were conducted following relevant guidelines and regulations to minimize animal suffering and distress in the laboratory. The Animal Care and Use Committee of Tohoku University (Permit number: 2019NiA-001) approved the animal experiments. The mice were monitored daily for their health during the full 4-week experiment duration. A humane end point was defined as a reduction of >25% of total body weight. The mice were euthanized through cervical dislocation, and death was verified through respiratory and cardiac arrest.

The mice were intraperitoneally immunized using 100 μg of the hCCR6p1-19C-KLH peptide with Imject Alum (Thermo Fisher Scientific, Inc.) to develop mAbs against hCCR6. The procedure included three additional immunizations, followed by a final booster intraperitoneal injection, 2 days before harvesting the spleen cells. PEG1500 (Roche Diagnostics, Indianapolis, IN) was used to subsequently fuse harvested spleen cells with P3U1 cells. Afterward, hybridomas were grown in the RPMI-1640 medium, supplemented with 10% heat-inactivated FBS, 100 units/mL penicillin, 100 $\mu\text{g}/\text{mL}$ streptomycin, 0.25 $\mu\text{g}/\text{mL}$ amphotericin B, and hypoxanthine/aminopterin/thymidine (HAT) for the selection (Thermo Fisher Scientific, Inc.).

Enzyme-linked immunosorbent assay (ELISA) with the hCCR6p1-19C peptide was used to subsequently screen supernatants, followed by flow cytometry, using CHO/hCCR6, CHO-K1, HepG2, and HuH-7.

Enzyme-linked immunosorbent assay

The hCCR6p1-19C peptide was immobilized on Nunc MaxiSorp 96-well immunoplates (Thermo Fisher Scientific, Inc.) at a concentration of 1 $\mu\text{g}/\text{mL}$ for 30 minutes at 37°C. After washing with phosphate-buffered saline (PBS) containing 0.05% Tween20 (PBST; Nacalai Tesque, Inc.), wells were blocked with 1% bovine serum albumin (BSA)-containing PBST for 30 minutes at 37°C. The plates were incubated with hybridoma supernatants, followed by peroxidase-conjugated anti-mouse immunoglobulins (1:2,000 diluted; Agilent Technologies, Inc., Santa Clara, CA). Enzymatic reactions were performed using the ELISA

POD Substrate TMB Kit (Nacalai Tesque, Inc.). Optical density was measured at 655 nm using an iMark Microplate Reader (Bio-Rad Laboratories, Inc., Berkeley, CA).

Flow cytometric analysis

Cells were harvested after a brief exposure to 0.25% trypsin and 1 mM of ethylenediaminetetraacetic acid (Nacalai Tesque, Inc.). They were washed with 0.1% BSA in PBS and treated with 1, 0.1, 0.01, or 0.001 $\mu\text{g}/\text{mL}$ of anti-hCCR6 mAbs for 30 minutes at 4°C. Then, the cells were treated with Alexa Fluor 488-conjugated anti-mouse IgG (1:2,000). Fluorescence data were obtained using SA3800 Cell Analyzer (Sony Corp.).

Determination of K_D by flow cytometry

The CHO/hCCR6, HepG2, and HuH-7 cells were suspended in 100 μL of serially diluted anti-hCCR6 mAbs, and then, 50 μL of Alexa Fluor 488-conjugated anti-mouse IgG (1:200) was added. The BD FACSLyric (BD Biosciences, Franklin Lakes) was used to obtain fluorescence data. The K_D was calculated by fitting saturation binding curves to built-in, one-site binding models in GraphPad Prism 6 (GraphPad Software, Inc., La Jolla).

Immunohistochemical analysis using $C_6\text{Mab-19}$ against non-Hodgkin lymphoma tissue

A formalin-fixed paraffin-embedded (FFPE) section of lymph nodes from non-Hodgkin lymphoma patients (Catalog No. T2235161-1; BioChain Institute, Inc., Newark, CA) was autoclaved in citrate buffer (pH6.0; Nichirei Biosciences, Inc., Tokyo, Japan) for 20 minutes. Then, the section was then incubated with 0.3 $\mu\text{g}/\text{mL}$ of $C_6\text{Mab-19}$ for 1 hour at room temperature, followed by treatment of an Envision+ kit (Agilent Technologies, Inc.) for 30 minutes at room temperature. The color was developed using 3,3'-diaminobenzidine tetrahydrochloride (DAB; Agilent Technologies, Inc.) for 1 minute at room temperature, and the section was then counterstained with hematoxylin (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan).

Results

Establishment of anti-hCCR6 mAbs using the peptide immunization method

In this study, we employed the peptide immunization method to develop anti-hCCR6 mAbs (Fig. 1). Two mice were immunized with hCCR6p1-19C-KLH (Fig. 1A). The spleens of the mice were excised, and splenocytes were subsequently fused with myeloma cells (P3U1) using PEG1500 (Fig. 1B). The produced hybridomas were seeded into 96-well plates and cultivated in HAT-contained medium for 7 days.

Hybridoma culture supernatants, which were positive for the hCCR6p1-19C peptide, were screened using ELISA (Fig. 1C). The first screening approach identified signals from hCCR6p1-19C in 134 out of 958 wells (14.0%). We selected the top 86 positive wells for the second screening. The second screening procedure using flow cytometry identified signals against CHO/hCCR6 cells in 49 out of the 86 hybridoma supernatants (57.0%). They did not react with parental CHO-

K1 cells. An anti-hCCR6 mAb, $C_6\text{Mab-19}$ (mouse IgG₁, kappa), was finally established after the single-cell cloning of hybridoma by limiting dilution and several additional flow cytometric screenings (Fig. 1D).

Flow cytometric analysis of $C_6\text{Mab-19}$ to hCCR6-expressing cells

We conducted flow cytometry using $C_6\text{Mab-19}$ and a commercially available anti-hCCR6 mAb (clone G034E3) against CHO/hCCR6, CHO-K1, HepG2, and HuH-7 cells. We selected HepG2 and HuH-7 cells for endogenously hCCR6-expressing cells because CCR6 expression was elevated in the human hepatocellular carcinoma cells in previous study.³² $C_6\text{Mab-19}$ dose dependently recognized CHO/hCCR6 cells at 1, 0.1, 0.01, and 0.001 $\mu\text{g}/\text{mL}$ (Fig. 2A), while $C_6\text{Mab-19}$ did not react to CHO-K1 parental cells even at 1 $\mu\text{g}/\text{mL}$ (Fig. 2B). Additionally, $C_6\text{Mab-19}$ possesses higher sensitivity than G034E3, especially at 0.1 and 0.01 $\mu\text{g}/\text{mL}$, although G034E3 recognized CHO/hCCR6 in a dose-dependent manner at 1, 0.1, 0.01, and 0.001 $\mu\text{g}/\text{mL}$ (Fig. 2A, C). Moreover, $C_6\text{Mab-19}$ also reacted to endogenously hCCR6-expressing cell lines, such as HepG2 and HuH-7 (Fig. 2D, E). $C_6\text{Mab-19}$ reacted to HepG2 at 1 and 0.1 $\mu\text{g}/\text{mL}$ (Fig. 2D) and reacted to HuH-7 at a low concentration of 0.01 $\mu\text{g}/\text{mL}$ (Fig. 2E). These results suggest that $C_6\text{Mab-19}$ is applicable for detecting exogenously and endogenously expressed hCCR6 using flow cytometry.

Determination of the K_D for anti-hCCR6 mAbs against hCCR6-expressing cells by flow cytometry

We performed the kinetic analysis using flow cytometry to assess the K_D of anti-hCCR6 mAbs, such as $C_6\text{Mab-19}$ and G034E3, for hCCR6-expressing cells. The K_D of $C_6\text{Mab-19}$ for CHO/hCCR6 cells was 3.0×10^{-10} M, as shown in Figure 3A. In contrast, the K_D of G034E3 for CHO/hCCR6 cells was 1.5×10^{-9} M (Fig. 3B), indicating a much higher binding affinity of $C_6\text{Mab-19}$ than that of G034E3.

Next, we analyzed the K_D of $C_6\text{Mab-19}$ against endogenously hCCR6-expressing HuH-7 and HepG2 cells. The K_D of $C_6\text{Mab-19}$ for HuH-7 and HepG2 was determined to be 1.8×10^{-10} M and 6.9×10^{-10} M, respectively (Fig. 3C, D). Therefore, $C_6\text{Mab-19}$ possesses a high affinity for exogenously and endogenously hCCR6-expressing cells.

Immunohistochemical analysis using $C_6\text{Mab-19}$ against non-Hodgkin lymphoma tissues

We stained an FFPE lymph node tissue from patients with non-Hodgkin lymphoma using $C_6\text{Mab-19}$ to investigate the use of $C_6\text{Mab-19}$ for immunohistochemical analysis. $C_6\text{Mab-19}$ strongly stained non-Hodgkin lymphoma cells (Fig. 4A, B). In contrast, no staining was observed without primary mAb (Fig. 4C, D). Therefore, $C_6\text{Mab-19}$ is useful for immunohistochemical staining of FFPE sections.

Discussion

We have previously established various mAbs against membrane proteins using the Cell-Based Immunization and Screening (CBIS) method. The CBIS method is a valid technique to develop mAbs against complex structured antigens, including GPCRs. Using the CBIS method, we




FIG. 1. A schematic illustration of the production flow of anti-hCCR6 mAbs. **(A)** The BALB/c mice were immunized with the hCCR6 peptide via the intraperitoneal route. **(B)** Hybridoma was produced by the fusion of P3U1 and mice splenocytes cells. **(C)** The first screening of hybridomas was conducted by ELISA against hCCR6p1–19C peptide, followed by flow cytometric secondary screening using hCCR6-expressing cells and parental cells. **(D)** Anti-hCCR6 mAbs were established after cloning via limiting dilution and additional analysis. ELISA, enzyme-linked immunosorbent assay; hCCR6, human CC chemokine receptor 6; mAbs, monoclonal antibodies.

successfully developed an anti-human CCR9 mAb (C₉Mab-1) and confirmed that its epitope is located at the N-terminus of human CCR9.^{30,33} By identifying the epitope of C₉Mab-1, we found that the N-terminal region of chemokine receptors is highly antigenic and have successfully developed mAbs against various mouse or human chemokine receptors, including CCR2,^{19,29} CCR3,^{20,23} CCR4,²⁴ CCR6,²⁵ CCR9,^{27,31} and CXCR6,²⁸ by using peptides in the N-terminal region as immunogens. Likewise, we successfully established a novel mAb against hCCR6 (clone C₆Mab-19), which is able to

recognize exogenous and endogenous hCCR6 by flow cytometry and immunohistochemistry.

The N-terminus of GPCRs, including CCR6, CCR9, and CXCR6, is identified as the ligand-binding domain.^{34–37} An anti-hCCR6 mAb (clone 1C6) can block the CCL20-CCR6 interaction and inhibit hCCR6-expressing cells' chemotaxis.³⁶ The epitope of 1C6 was determined as the N-terminus of hCCR6.³⁶ CCL20 binds to all three extracellular domains of CCR6 and N-terminal residues from Tyr27 to Leu38.⁷ Conversely, N-terminal residues before Tyr27 in CCR6 may




FIG. 2. The analysis of reactivity of $C_6\text{Mab-19}$ by flow cytometry. CHO/hCCR6 (A) or CHO-K1 (B) cells were treated with 0.001, 0.01, 0.1, and 1 $\mu\text{g/mL}$ of $C_6\text{Mab-19}$, followed by treatment with Alexa Fluor 488-conjugated anti-mouse IgG (red line). (C) CHO/hCCR6 cells were treated with 0.001, 0.01, 0.1, and 1 $\mu\text{g/mL}$ of G034E3, which is a commercially available anti-hCCR6 mAb from BioLegend, followed by treatment with Alexa Fluor 488-conjugated anti-mouse IgG (red line). HepG2 (D) or HuH-7 (E) cells were treated with 0.001, 0.01, 0.1, and 1 $\mu\text{g/mL}$ of $C_6\text{Mab-19}$, followed by treatment with Alexa Fluor 488-conjugated anti-mouse IgG (red line). The black line shows the negative control (blocking buffer). CHO, Chinese hamster ovary.




FIG. 3. The determination of the K_D of anti-hCCR6 mAbs against hCCR6-expressing cells. CHO/hCCR6 (A, B), HuH-7 (C), and HepG2 (D) cells were suspended in serially diluted C₆Mab-19 or G034E3 at the indicated concentrations. Then, cells were treated with Alexa Fluor 488-conjugated anti-mouse IgG. Fluorescence data were subsequently obtained using BD FACSLytic, followed by the calculation of the dissociation constant (K_D) by GraphPad Prism 6.




FIG. 4. The immunohistochemical staining using C₆Mab-19 against non-Hodgkin lymphoma tissues. After antigen retrieval, serial tissue sections of non-Hodgkin lymphoma (Catalog number: T2235161-1; BioChain) were incubated with 0.3 $\mu\text{g}/\text{mL}$ of C₆Mab-19 (A, B), or without primary antibody (C, D) followed by treatment with the Envision+ kit. The color development was performed using DAB, and the sections were counterstained with hematoxylin. The scale bar indicated 100 μm . DAB, 3,3'-diaminobenzidine tetrahydrochloride.

not contribute to the interaction with CCL20.⁷ The second extracellular domain and N-terminal residues from Tyr27 to Leu38 in CCR6 are the central sites of interaction in CCR6. The epitope of C₆Mab-19 is presented in the 1st to 19th amino acids of hCCR6; therefore, C₆Mab-19 might not possess the neutralizing activity for CCL20. We try to establish the mAbs, which have the neutralizing activity for CCL20 by applying the peptide immunization method, in the future studies.

CCL20/CCR6 has been reported to be involved in many inflammatory and autoimmune diseases, and the CCL20/CCR6 axis has been a therapeutic target.⁴ Small molecule inhibitors for CCL20/CCR6, such as compound 1b and MR120 are expected to ameliorate symptoms of chronic colitis in preclinical models.^{10,38} Interestingly, a recent report revealed that the high prevalence of CCR6 in patients with acute myocardial infarction makes it an independent biomarker and correlates with the severity.³⁹ Lipid-lowering drugs ameliorate psoriasis by suppressing inflammatory responses that also involve CCL20/CCR6.⁴⁰

Furthermore, oral squamous cell carcinoma patients with drinking and smoking habits significantly increased the percentage of CD4⁺ T cells, which express CD45RO and CCR6 in the blood and tumor microenvironment compared with the patients who do not drink or smoke.⁴¹ Hence, CCR6 is a potent immunoregulatory factor in the formation of pathologies. Therefore, CCR6 is an attractive target to help patients with various inflammatory diseases. C₆Mab-19 can be used in flow cytometry (Fig. 2) and immunohistochemistry (Fig. 4); thus, it may contribute to the diagnosis of such diseases. Additionally, C₆Mab-19 may treat such inflammatory diseases, although further validation is needed.

CCL20, which is secreted into tumor tissues, recruits immunosuppressive CCR6-expressing cells, including Treg cells.¹³ CCL20/CCR6-guided circulating Tregs infiltrate into tumors, resulting in tumor progression and poor prognosis in patients with hepatocellular carcinoma.¹⁴ Hence, cancer treatment strategies using CCR6-expressing CAR-T cells were designed.^{18,42} We could detect CCR6-expressing CAR-T cells in blood or tissues using flow cytometry or immunohistochemistry because C₆Mab-19 possesses extremely high binding affinity against hCCR6.

Author Disclosure Statement

The authors have no conflict of interest.

Funding Information

This research was supported in part by Japan Agency for Medical Research and Development (AMED) under Grant Numbers: JP22ama121008 (to Y.K.), JP22am0401013 (to Y.K.), JP22bm1004001 (to Y.K.), JP22ck0106730 (to Y.K.), and JP21am0101078 (to Y.K.), and by Japan Society for the Promotion of Science (JSPS) Grants-in-Aid for Scientific Research (KAKENHI) grant nos. 21K20789 (to T.T.), 22K06995 (to H.S.), 21K07168 (to M.K.K.), and 22K07224 (to Y.K.).

References

- Zlotnik A, Yoshie O. The chemokine superfamily revisited. *Immunity* 2012;36:705–716; doi: 10.1016/j.immuni.2012.05.008
- Hughes CE, Nibbs RJB. A guide to chemokines and their receptors. *FEBS J* 2018;285:2944–2971; doi: 10.1111/febs.14466
- Nagarsheth N, Wicha MS, Zou W. Chemokines in the cancer microenvironment and their relevance in cancer immunotherapy. *Nat Rev Immunol* 2017;17:559–572; doi: 10.1038/nri.2017.49
- Ranasinghe R, Eri R. Modulation of the CCR6-CCL20 axis: A potential therapeutic target in inflammation and cancer. *Medicina (Kaunas)* 2018;54(5):88; doi: 10.3390/medicina54050088
- Schöneberg T, Liebscher I. Mutations in G protein-coupled receptors: Mechanisms, pathophysiology and potential therapeutic approaches. *Pharmacol Rev* 2021;73:89–119; doi: 10.1124/pharmrev.120.000011
- Baba M, Imai T, Nishimura M, et al. Identification of CCR6, the specific receptor for a novel lymphocyte-directed CC chemokine LARC. *J Biol Chem* 1997;272:14893–14898; doi: 10.1074/jbc.272.23.14893
- Wasilko DJ, Johnson ZL, Ammirati M, et al. Structural basis for chemokine receptor CCR6 activation by the endogenous protein ligand CCL20. *Nat Commun* 2020;11:3031; doi: 10.1038/s41467-020-16820-6
- Ye C, Guo X, Wu J, et al. CCL20/CCR6 mediated macrophage activation and polarization can promote adenoid epithelial inflammation in adenoid hypertrophy. *J Inflamm Res* 2022;15:6843–6855; doi: 10.2147/jir.S390210
- Tanaka R, Ichimura Y, Kubota N, et al. The role of PD-L1 on langerhans cells in the regulation of psoriasis. *J Invest Dermatol* 2022;142:3167–3174.e3169; doi: 10.1016/j.jid.2022.06.006
- Martina MG, Giorgio C, Allodi M, et al. Discovery of small-molecules targeting the CCL20/CCR6 axis as first-in-class inhibitors for inflammatory bowel diseases. *Eur J Med Chem* 2022;243:114703; doi: 10.1016/j.ejmech.2022.114703
- Kadomoto S, Izumi K, Mizokami A. The CCL20-CCR6 axis in cancer progression. *Int J Mol Sci* 2020;21(15):5186; doi: 10.3390/ijms21155186
- Lee JJ, Kao KC, Chiu YL, et al. Enrichment of human CCR6(+) regulatory T cells with superior suppressive activity in oral cancer. *J Immunol* 2017;199:467–476; doi: 10.4049/jimmunol.1601815
- Rutihinda C, Haroun R, Saidi NE, et al. Inhibition of the CCR6-CCL20 axis prevents regulatory T cell recruitment and sensitizes head and neck squamous cell carcinoma to radiation therapy. *Cancer Immunol Immunother* 2023;72(5):1089–1102; doi: 10.1007/s00262-022-03313-2
- Chen KJ, Lin SZ, Zhou L, et al. Selective recruitment of regulatory T cell through CCR6-CCL20 in hepatocellular carcinoma fosters tumor progression and predicts poor prognosis. *PLoS One* 2011;6:e24671; doi: 10.1371/journal.pone.0024671
- Walch-Rückheim B, Mavrova R, Henning M, et al. Stromal fibroblasts induce CCL20 through IL6/C/EBPβ to support the recruitment of Th17 cells during cervical cancer progression. *Cancer Res* 2015;75:5248–5259; doi: 10.1158/0008-5472.Can-15-0732
- Kadomoto S, Izumi K, Hiratsuka K, et al. Tumor-associated macrophages induce migration of renal cell carcinoma cells via activation of the CCL20-CCR6 axis. *Cancers (Basel)* 2019;12(1):89; doi: 10.3390/cancers12010089
- Liu W, Wang W, Wang X, et al. Cisplatin-stimulated macrophages promote ovarian cancer migration via the CCL20-CCR6 axis. *Cancer Lett* 2020;472:59–69; doi: 10.1016/j.canlet.2019.12.024

18. Jin L, Cao L, Zhu Y, et al. Enhance anti-lung tumor efficacy of chimeric antigen receptor-T cells by ectopic expression of C-C motif chemokine receptor 6. *Sci Bull (Beijing)* 2021; 66:803–812; doi: 10.1016/j.scib.2020.12.027
19. Tanaka T, Li G, Asano T, et al. Development of a novel anti-mouse CCR2 monoclonal antibody (C(2)Mab-6) by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:80–86; doi: 10.1089/mab.2021.0063
20. Asano T, Nanamiya R, Takei J, et al. Development of anti-mouse CC chemokine receptor 3 monoclonal antibodies for flow cytometry. *Monoclon Antib Immunodiagn Immunother* 2021;40:107–112; doi: 10.1089/mab.2021.0009
21. Saito M, Harigae Y, Li G, et al. C(3)Mab-2: An anti-mouse CCR3 monoclonal antibody for immunocytochemistry. *Monoclon Antib Immunodiagn Immunother* 2022;41:45–49; doi: 10.1089/mab.2021.0050
22. Asano T, Suzuki H, Tanaka T, et al. C(3)Mab-3: A monoclonal antibody for mouse CC chemokine receptor 3 for flow cytometry. *Monoclon Antib Immunodiagn Immunother* 2022;41:74–79; doi: 10.1089/mab.2021.0062
23. Asano T, Suzuki H, Goto N, et al. Establishment of novel anti-mouse CCR3 monoclonal antibodies (C(3)Mab-6 and C(3)Mab-7) by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:94–100; doi: 10.1089/mab.2021.0065
24. Takei J, Suzuki H, Asano T, et al. Development of a novel anti-mouse CCR4 monoclonal antibody (C(4)Mab-1) by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:87–93; doi: 10.1089/mab.2021.0064
25. Asano T, Tanaka T, Suzuki H, et al. Development of a novel anti-mouse CCR6 monoclonal antibody (C(6)Mab-13) by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:343–349; doi: 10.1089/mab.2022.0021
26. Tanaka T, Nanamiya R, Takei J, et al. Development of anti-mouse CC chemokine receptor 8 monoclonal antibodies for flow cytometry. *Monoclon Antib Immunodiagn Immunother* 2021;40:65–70; doi: 10.1089/mab.2021.0005
27. Kobayashi H, Asano T, Suzuki H, et al. Establishment of a sensitive monoclonal antibody against mouse CCR9 (C(9)Mab-24) for flow cytometry. *Monoclon Antib Immunodiagn Immunother* 2023;42(1):15–21; doi: 10.1089/mab.2022.0032
28. Kitamura K, Suzuki H, Kaneko MK, et al. Cx(6)Mab-1: A novel anti-mouse CXCR6 monoclonal antibody established by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:133–141; doi: 10.1089/mab.2022.0010
29. Tanaka T, Li G, Saito M, et al. Development of an anti-human CCR2 monoclonal antibody (C(2)Mab-9) by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:188–193; doi: 10.1089/mab.2022.0001
30. Nanamiya R, Takei J, Asano T, et al. Development of anti-human CC chemokine receptor 9 monoclonal antibodies for flow cytometry. *Monoclon Antib Immunodiagn Immunother* 2021;40:101–106; doi: 10.1089/mab.2021.0007
31. Tanaka T, Suzuki H, Isoda Y, et al. Development of a sensitive anti-human CCR9 monoclonal antibody (C(9)Mab-11) by N-terminal peptide immunization. *Monoclon Antib Immunodiagn Immunother* 2022;41:303–310; doi: 10.1089/mab.2022.0027
32. Fujii H, Itoh Y, Yamaguchi K, et al. Chemokine CCL20 enhances the growth of HuH7 cells via phosphorylation of p44/42 MAPK in vitro. *Biochem Biophys Res Commun* 2004;322:1052–1058; doi: 10.1016/j.bbrc.2004.07.207
33. Takei J, Asano T, Li G, et al. Epitope mapping of an anti-human CCR9 monoclonal antibody (C(9)Mab-1) using enzyme-linked immunosorbent assay. *Monoclon Antib Immunodiagn Immunother* 2021;40:239–242; doi: 10.1089/mab.2021.0037
34. Chamorro S, Vela M, Franco-Villanueva A, et al. Anti-tumor effects of a monoclonal antibody to human CCR9 in leukemia cell xenografts. *MAbs* 2014;6:1000–1012; doi: 10.4161/mabs.29063
35. Chain B, Arnold J, Akthar S, et al. A linear epitope in the N-terminal domain of CCR5 and its interaction with antibody. *PLoS One* 2015;10:e0128381; doi: 10.1371/journal.pone.0128381
36. Gómez-Melero S, García-Maceira FI, García-Maceira T, et al. Amino terminal recognition by a CCR6 chemokine receptor antibody blocks CCL20 signaling and IL-17 expression via β -arrestin. *BMC Biotechnol* 2021;21:41; doi: 10.1186/s12896-021-00699-2
37. Aguilera-Durán G, Romo-Mancillas A. Behavior of chemokine receptor 6 (CXCR6) in complex with CXCL16 soluble form chemokine by molecular dynamic simulations: General protein–ligand interaction model and 3D-QSAR studies of synthetic antagonists. *Life (Basel)* 2021; 11(4):346; doi: 10.3390/life11040346
38. Allodi M, Giorgio C, Incerti M, et al. Probing the effects of MR120 in preclinical chronic colitis: A first-in-class anti-IBD agent targeting the CCL20/CCR6 axis. *Eur J Pharmacol* 2023;945:175613; doi: 10.1016/j.ejphar.2023.175613
39. Abdelrahman AMN, Mohamed AE, Bendary AM, et al. Assessment of circulating CCR6 level in acute myocardial infarction and its association with disease severity. *Cardiovasc Hematol Agents Med Chem* 2023;21(3):223–231; doi: 10.2174/1871525721666230120160717
40. Wang J, Zhang S, Xing M, et al. Current evidence on the role of lipid lowering drugs in the treatment of psoriasis. *Front Med (Lausanne)* 2022;9:900916; doi: 10.3389/fmed.2022.900916
41. Rochefort J, Karagiannidis I, Baillou C, et al. Defining biomarkers in oral cancer according to smoking and drinking status. *Front Oncol* 2022;12:1068979; doi: 10.3389/fonc.2022.1068979
42. Zhong C, Chen J. CAR-T cell engineering with CCR6 exhibits superior anti-solid tumor efficacy. *Sci Bull (Beijing)* 2021;66:755–756; doi: 10.1016/j.scib.2020.12.021

Address correspondence to:

Yukinari Kato
 Department of Antibody Drug Development
 Tohoku University Graduate School of Medicine
 2-1, Seiryomachi, Aoba-ku
 Sendai 980-8575
 Miyagi
 Japan

E-mail: yukinari.kato.e6@tohoku.ac.jp

Received: March 25, 2023

Accepted: June 17, 2023