

COMMUNICATION

C₈Mab-21: A novel anti-human CCR8 monoclonal antibody for flow cytometry

Haruto Yamamoto[†], Yu Kaneko[†], Tomohiro Tanaka[†], Guanjie Li^{ID},
Hiroyuki Suzuki^{*ID}, Mika K. Kaneko^{ID}, and Yukinari Kato^{*ID}

Department of Antibody Drug Development, Tohoku University Graduate School of Medicine, 2-1, Seiryō-machi, Aoba-ku, Sendai, Miyagi, Japan

(This article belongs to the *Special Issue: Immunomodulation and Antitumor Strategies in the Tumor Microenvironment*)

Abstract

C-C motif chemokine receptor-8 (CCR8) belongs to class A of G protein-coupled receptors. CCR8 interacts with the specific chemokine ligand CCL1/I-309 in humans, which is produced by various cells, including tumor-associated macrophages and regulatory T cells (Treg). CCR8 is highly expressed on Treg and T-helper 2 cells recruited to the inflammation site and is implicated in allergy, asthma, and cancer progression. CCR8⁺Treg cells have been suggested an important regulator in the immunosuppressive tumor microenvironment. Therefore, it has been proposed for use in the development of sensitive monoclonal antibodies targeting CCR8. This study developed a specific mAb for human CCR8 (hCCR8), which is useful for flow cytometry by employing the Cell-Based Immunization and Screening (CBIS) method. The established anti-hCCR8 mAb (C₈Mab-21; mouse IgM, kappa) demonstrated reactivity with hCCR8-overexpressed Chinese hamster ovary-K1 (CHO/hCCR8) cells, TALL-1 (human adult acute T-lymphoblastic leukemia), CCRF-HSB2 (human T-lymphoblastic leukemia), and natural killer cells expressing endogenous hCCR8, as confirmed by flow cytometry. Furthermore, EC₅₀ values of C₈Mab-21 for CHO/hCCR8 and TALL-1 were determined as 6.5×10^{-8} M and 2.0×10^{-8} M, respectively. C₈Mab-21, established by the CBIS method, provides a useful tool for analyzing the hCCR8-related biological response using flow cytometry.

Keywords: CCR8; CBIS method; Monoclonal antibody; Flow cytometry

[†]These authors contributed equally to this work.

***Corresponding authors:**

Hiroyuki Suzuki
(hiroyuki.suzuki.b4@tohoku.ac.jp)
Yukinari Kato
(yukinari.kato.e6@tohoku.ac.jp)

Citation: Yamamoto H, Kaneko Y, Tanaka T, *et al.* C₈Mab-21: A novel anti-human CCR8 monoclonal antibody for flow cytometry. *Microbes & Immunity*. 2025; 2(2):126-136.
doi: 10.36922/mi.4661

Received: August 26, 2024

Revised: November 13, 2024

Accepted: December 2, 2024

Published online: December 16, 2024

Copyright: © 2024 Author(s).

This is an Open-Access article distributed under the terms of the Creative Commons Attribution License, permitting distribution, and reproduction in any medium, which provided that the original work is properly cited.

Publisher's Note: AccScience Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

1. Introduction

Targeting immune checkpoint has become an effective and powerful strategy for cancer therapy.¹⁻⁴ In particular, the development of antibody drugs targeting immune checkpoint molecules, such as programmed-cell death-1 (PD-1), cytotoxic T lymphocyte antigen 4 (CTLA-4), and PD-1 ligand 1 (PD-L1), has achieved remarkable therapeutic results.⁵⁻⁷ PD-1 inhibits the excessive activation of conventional T cells by suppressing costimulatory signaling and renders them dysfunctional or exhausted.⁸ PD-1 and CTLA-4 are also expressed in regulatory T cells (Treg), one of the immunosuppresses in the tumor microenvironment (TME).⁹ Inhibition of

these molecules could potentiate the activation and immunosuppressive function of Treg.¹⁰⁻¹²

Treg is defined as CD4⁺T cell that expresses CD25 and FOXP3, playing a role in maintaining self-tolerance to prevent excessive immune responses and autoimmune diseases.^{13,14} Treg suppresses the effector functions of T cells through the secretion of immunosuppressive cytokines, such as interleukin-10 (IL-10), transforming growth factor- β , and cytotoxic granzyme/perforin.¹⁴⁻¹⁸ Intratumoral Treg suppresses antitumor T cell responses and thus resists the effects of immune checkpoint inhibitor therapy.^{19,20} Antibodies against T cell immunoreceptors with Ig and ITIM domains (TIGIT), one of the immune checkpoint molecules, have shown to improve the effectiveness of PD-L1 antibodies by suppressing Treg.²¹ Therefore, the development of immunotherapy targeting Treg is expected.^{11,22}

Intratumoral Treg expresses high levels of C-C motif chemokine receptor-8 (CCR8).⁴ In addition, CCR8-expressing Tregs exhibit increased expression of CD25 and FOXP3 compared to CCR8-negative Tregs,²³ indicating their potent immunosuppressive functions. The CCR8-expressing Treg is known to be correlated with poor prognosis in some cancer patients.^{24,25} Thus, CCR8 is emerging as an attractive target for the next cancer immunotherapy.²⁶ Several anti-CCR8 drugs, including S-531011,²⁷ IPG7236,²⁸ and SRF114²⁹ are undergoing clinical trials.

CCR8 is one of the seven transmembrane-spanning G protein-coupled receptors.⁴ Human CCR8 (hCCR8) is known to bind to five C-C chemokine ligands (CCLs): CCL1/I-309, CCL4, CCL16, CCL17, and CCL18.³⁰⁻³² CCR8 is upregulated not only in Treg but also in various cancers, including breast, non-small cell lung (NSCLC), bladder, and colorectal cancer.^{24,25} In bladder cancer, CCR8 mediates cell migration, invasion, and epithelial-mesenchymal transition by interacting with CCL18.³³ In addition, CCR8 and its specific ligand CCL1/I-309 regulate the immune system, which mediates the progression of diseases such as cancers by promoting migration and inhibiting apoptosis in Treg and lymphomas.^{34,35} Therefore, CCR8-targeting antibodies will contribute to the elucidation of pathological mechanisms, diagnosis, and therapy.^{27,36}

Using the Cell-Based Immunization and Screening (CBIS) method, we previously developed numerous monoclonal antibodies (mAbs) against chemokine receptors, including mouse CCR3,³⁷ mouse CCR8,³⁸ human CCR9,³⁹ and mouse C-X-C chemokine receptor type 4 (CXCR4).⁴⁰ In this study, we successfully developed an anti-hCCR8 mAb using the CBIS method, which is applicable to flow cytometry.

2. Materials and methods

2.1. Cell lines

LN229, Chinese hamster ovary (CHO)-K1, and P3X63Ag8U.1 (P3U1) cells were obtained from the American Type Culture Collection (Manassas, VA, USA). TALL-1 and CCRF-HSB2 cells were obtained from the Japanese Collection of Research Bioresources Cell Bank (Osaka, Japan). The human natural killer (NK) cells (donor lot. 4022602, purity >70%) were purchased from Takara Bio (Shiga, Japan). pCMV6neo-myc-DDK vector with hCCR8 (Accession No.: NM_005201) was purchased from OriGene Technologies, Inc. (Rockville, MD, USA). The plasmid was transfected into cell lines using the Neon transfection system (Thermo Fisher Scientific, Inc., Waltham, MA, USA). Subsequently, LN229 and CHO-K1 stably overexpressing hCCR8 with C-terminal myc-DDK tags (hereinafter described as LN229/hCCR8 and CHO/hCCR8, respectively) were established using a cell sorter (SH800; Sony Corp., Tokyo, Japan), following cultivation in a medium containing 0.5 mg/mL G418 (Nacalai Tesque, Inc., Kyoto, Japan).

CHO-K1, P3U1, CHO/hCCR8, TALL-1, and CCRF-HSB2 cells were cultured in a Roswell Park Memorial Institute (RPMI)-1640 medium (Nacalai Tesque, Inc., Kyoto, Japan) supplemented with 10% heat-inactivated fetal bovine serum (FBS, Thermo Fisher Scientific Inc., Waltham, MA, USA), 100 units/mL penicillin, 100 μ g/mL streptomycin, and 0.25 μ g/mL amphotericin B (Nacalai Tesque, Inc., Kyoto, Japan). LN229 and LN229/hCCR8 were cultured in a Dulbecco's Modified Eagle Medium (DMEM, Nacalai Tesque, Inc., Kyoto, Japan) supplemented with 10% heat-inactivated FBS (Thermo Fisher Scientific Inc., Waltham, MA, USA), 100 units/mL penicillin, 100 μ g/mL streptomycin, and 0.25 μ g/mL amphotericin B (Nacalai Tesque, Inc., Kyoto, Japan). All cells were cultured in a humidified incubator at 37°C with 5% CO₂ and 95% air.

2.2. Antibodies

The anti-human CD198 (CCR8) mAb (clones S19017D and L263G8) were purchased from BioLegend (San Diego, CA, USA). The Alexa Fluor 488-conjugated anti-mouse Immunoglobulin (Ig)G was purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA).

2.3. Hybridoma production

For developing anti-hCCR8 mAbs, two female 6-week-old BALB/c mice were immunized intraperitoneally with 1×10^8 cells of LN229/hCCR8. The immunogen was harvested after brief exposure to 1 mM ethylenediaminetetraacetic acid (EDTA; Nacalai Tesque, Inc., Kyoto, Japan). Imject

Alum (Thermo Fisher Scientific Inc., Waltham, MA, USA) was used as an adjuvant for the first immunization. Subsequently, three additional weekly injections of 1×10^8 cells of LN229/hCCR8 were administered without an adjuvant. A final booster immunization with 1×10^8 cells of LN229/hCCR8 was given intraperitoneally 2 days before harvesting splenocytes from the mice. Harvested splenocytes were then fused with P3U1 cells using polyethylene glycol 1500 (PEG1500; Roche Diagnostics, Indianapolis, IN, USA).

Hybridomas were cultured in RPMI-1640 medium supplemented as shown above and additional supplements included hypoxanthine, aminopterin, and thymidine (HAT; Thermo Fisher Scientific, Inc., Waltham, MA, USA), 5% Briclone (NICB, Dublin, Ireland), and 5 $\mu\text{g/mL}$ of Plasmocin (InvivoGen, San Diego, CA, USA). The hybridoma supernatants were screened by flow cytometry using CHO/hCCR8 and parental CHO-K1 cells. The cultured supernatant of C₈Mab-21-producing hybridomas was filtrated and purified using Capto L (Cytiva, Tokyo, Japan).

2.4. Flow cytometry

CHO-K1 and CHO/hCCR8 cells were harvested after brief exposure to 1 mM EDTA. Subsequently, CHO-K1, CHO/hCCR8, TALL-1, and CCRF-HSB2 cells were washed with 0.1% bovine serum albumin in phosphate-buffered saline and treated with primary mAbs for 30 min at 4°C. The cells were then treated with Alexa Fluor 488-conjugated anti-mouse IgG (1:1000) following the collection of fluorescence data, using the SA3800 Cell Analyzer and SA3800 software ver. 2.05 (Sony Corp, Tokyo, Japan).

2.5. Determination of the EC₅₀ by flow cytometry

CHO/hCCR8 and TALL-1 were suspended in 100 μL of serially diluted C₈Mab-21 (100 $\mu\text{g/mL}$ – 0.006 $\mu\text{g/mL}$), S19017D (10 $\mu\text{g/mL}$ – 0.0006 $\mu\text{g/mL}$ for CHO/hCCR8; 10 $\mu\text{g/mL}$, 2.5 $\mu\text{g/mL}$ – 0.0006 $\mu\text{g/mL}$ for TALL-1), or L263G8 (10 $\mu\text{g/mL}$ – 0.0006 $\mu\text{g/mL}$ for CHO/hCCR8; 0.625 $\mu\text{g/mL}$ – 0.0006 $\mu\text{g/mL}$ for TALL-1). Alexa Fluor 488-conjugated anti-mouse IgG (1:200) was then added. Fluorescence data were subsequently collected using the BD FACSLytic (BD Biosciences, Franklin Lakes, NJ, USA), and EC₅₀ values were calculated by fitting the binding isotherms into the built-in one-site binding model in GraphPad PRISM 6 (GraphPad Software, Inc., La Jolla, CA, USA).

3. Results

3.1. Establishment of anti-hCCR8 mAbs by the CBIS method


The CBIS method was employed using hCCR8-overexpressing cells to develop anti-hCCR8 mAbs. Anti-

hCCR8 mAbs-producing hybridoma screening was conducted using flow cytometry (Figure 1). Two mice were intraperitoneally immunized with LN229/hCCR8 weekly for a total of 5 times. Subsequently, hybridomas were seeded into 96-well plates, after which flow cytometric analysis was used to select CHO/hCCR8-reactive and CHO-K1-non-reactive supernatants of hybridomas. From 956 wells, only one (0.10%) yielded CHO/hCCR8-reactive supernatant. Finally, we established the clone C₈Mab-21 (mouse IgM, kappa) by limiting dilution and additional screening.


A Immunization of hCCR8-expressing Cells



B Production of hybridomas



C Screening by using flow cytometry



D Cloning of anti-hCCR8 mAb-producing hybridomas




Figure 1. A schematic procedure of anti-hCCR8 monoclonal antibodies production. The procedure of the CBIS method for antibody development. LN229/hCCR8 cells were immunized into two mice via intraperitoneal injection (A). Splenic cells harvested from mice were fused with P3U1 myeloma cells (B). The culture supernatants of hybridoma were screened by flow cytometry using CHO-K1 and CHO/hCCR8 (C). After limiting dilution of hybridomas and additional analysis, the C₈Mab-21 clone was finally established (D).

Abbreviations: i.p.: Intraperitoneal; CHO: Chinese hamster ovary; CBIS: Cell-based immunization and screening; hCCR8: Human C-C motif chemokine receptor-8.

3.2. Flow cytometric analysis

Flow cytometric analysis was conducted using purified C₈Mab-21 (Figure S1) and commercially available anti-human CD198 (CCR8) mAbs (clone S19017D and L263G8) against CHO-K1, CHO/hCCR8, TALL-1, CCRF-HSB2, and NK cells. The results showed that C₈Mab-21, S19017D, and L263G8 recognized CHO/hCCR8 in a dose-dependent manner (Figure 2A). Neither C₈Mab-21 nor L263G8 reacted with parental CHO-K1 cells, even at a concentration of 20 µg/mL. However, S19017D showed slight reactivity with CHO-K1 cells at concentrations of 20 µg/mL and 2 µg/mL (Figure 2B). For endogenously hCCR8-expressing cells, C₈Mab-21 recognized TALL-1, CCRF-HSB2, and NK cells at concentrations of 2 µg/mL and 20 µg/mL (Figure 3). S19017D reacted with TALL-1 and CCRF-HSB2 in a dose-dependent manner, even at a concentration of 0.02 µg/mL, but did not recognize NK cells, even at a concentration of 20 µg/mL (Figure 3). L263G8 reacted with TALL-1 and CCRF-HSB2 at concentration as low as 0.02 µg/mL, and with NK cells at concentrations of 2 µg/mL or higher (Figure 3). Thus, C₈Mab-21 could detect both exogenously and endogenously expressed hCCR8 in its native conformation using flow cytometry.

3.3. Titration of anti-CCR8 mAbs on hCCR8-overexpressed and endogenously hCCR8-expressing cell lines

The titration of C₈Mab-21, S19017D, and L263G8 was assessed with exogenously hCCR8-expressed CHO/hCCR8 using flow cytometry. The results showed that the EC₅₀ values of C₈Mab-21, S19017D, and L263G8 for CHO/hCCR8 are 6.5×10^{-8} M, 2.6×10^{-9} M, and 1.2×10^{-9} M, respectively (Figure 4). The histogram of C₈Mab-21 is shown in Figure S2. These results indicate that C₈Mab-21 possesses a moderate affinity for CHO/hCCR8 cells.

The titration of C₈Mab-21, S19017D, and L263G8 was analyzed with endogenously hCCR8-expressing TALL-1 using flow cytometry. The EC₅₀ values of C₈Mab-21, S19017D, and L263G8 for TALL-1 are 2.0×10^{-8} M, 4.6×10^{-10} M, and 7.8×10^{-11} M, respectively (Figure 5). These results showed that C₈Mab-21 possesses a moderate affinity for endogenously expressed hCCR8 in TALL-1 leukemia cells.

Further investigation was conducted to explore other applications, such as immunohistochemistry. However, hCCR8 could not be detected by immunohistochemistry using cell blocks of CHO/hCCR8 (Figure S3).

4. Discussion

Chemokine receptors are focused as targets for many diseases, including inflammatory disorders and cancers.^{41,42}

The receptors transmit signals to intracellular molecules regarding extracellular conditions and govern broad cellular dynamics, such as proliferation, homeostasis, migration, and motility of the cells.^{41,43} Although the therapeutic drugs including mAbs have been developed, significant challenges remain. These challenges include the complexity of the structure, the small area of epitope regions, and the difficulty in purifying these protein as immunogens.^{42,44}

Unlike protein purifications, the preparation of antigens using the CBIS method is relatively simple. Furthermore, the CBIS method allows for the retention of the antigens' structure, including modifications such as glycosylation and folding. We have successfully developed multiple mAbs using the CBIS method, targeting human epidermal growth factor receptor 1 (HER1; EGFR),⁴⁵ HER3,⁴⁶ trophoblast cell surface antigen 2,⁴⁷ CD44,⁴⁸ and podoplanin.⁴⁹ Furthermore, some of the mAbs developed by the CBIS method exhibited cancer specificity by recognizing unique cancer-specific epitopes.^{50,51} Therefore, the CBIS method is one of the efficient and useful tactics for generating diverse antibodies targeting membrane proteins. Further investigations are required to determine the epitope of C₈Mab-21.

In the immunosuppressive TME, CD8⁺T cells are exhausted along with the induction of CCR8⁺Treg. The infiltration of CCR8⁺Treg has been shown to associate with high thymocyte selection associated high mobility group box (TOX), an exhaustion marker in CD8⁺T cells, in some types of cancer patients.^{52,53} Elimination of CCR8⁺Treg using antibodies is expected to advance the treatment of these cancer patients. Targeting CCR8 may offer more specific antitumor activity than other approaches aimed at Treg removal.^{53,54} In mice, CCR8⁺T cell depletion therapy using anti-CCR8 mAbs induces tumor-specific immune responses without triggering autoimmune responses or immune reactions in the TME.³⁶ Since C₈Mab-21 recognizes cell surface hCCR8, we plan to investigate its potential function against Treg, such as detection and interfering effects, in future studies. Furthermore, antibody-dependent cellular cytotoxicity (ADCC) activity and complement-dependent cytotoxicity have previously been enhanced by modifying isotypes and defucosylating mAbs.^{55,56} C₈Mab-21 is a mouse IgM, which lacks ADCC activity. Although the crosslinking property of IgM is lost, it will be converted into a mouse IgG_{2a} version to evaluate the effect of antitumor activities in xenograft models. We successfully cloned and determined the complementarity-determining regions of C₈Mab-21 (Figure S4).

Interestingly, a correlation between cancer-associated fibroblasts (CAFs) and CCR8 has been found from




Figure 2. Flow cytometric analysis of anti-hCCR8 monoclonal antibodies against CHO/hCCR8 and CHO-K1. CHO/hCCR8 (A) and CHO-K1 (B) cells were treated with 0.02 – 20 $\mu\text{g/mL}$ of C₈Mab-21, S19017D, and L263G8 (red line), followed by treatment with Alexa Fluor 488-conjugated anti-mouse Immunoglobulin G. Fluorescence data were collected using the SA3800 Cell Analyzer. The black line represents the control group (no primary antibody treatment).

Abbreviations: CHO: Chinese hamster ovary; hCCR8: Human C-C motif chemokine receptor-8.




Figure 3. Flow cytometric analysis of anti-hCCR8 monoclonal antibodies against endogenously hCCR8-expressing cells. TALL-1 (A), CCRF-HSB2 (B), and NK cells (C) were treated with 0.02 – 20 µg/mL of C₈Mab-21, S19017D, and L263G8 (red line), followed by treatment with Alexa Fluor 488-conjugated anti-mouse Immunoglobulin G. Fluorescence data were collected using the SA3800 Cell Analyzer. The black line represents the control group (no primary antibody treatment).

Abbreviations: NK cells: Natural killer cells; hCCR8: Human C-C motif chemokine receptor-8.




Figure 4. The analysis of the binding affinity of anti-hCCR8 monoclonal antibodies for CHO/hCCR8. CHO/hCCR8 cells were suspended in 100 μ L of serially diluted C₈Mab-21 (100 μ g/mL – 0.006 μ g/mL) (A), S19017D (10 μ g/mL – 0.0006 μ g/mL) (B), or L263G8 (10 μ g/mL – 0.0006 μ g/mL) (C). The cells were then treated with Alexa Fluor 488-conjugated anti-mouse immunoglobulin G. Fluorescence data were subsequently collected using the BD FACSLyric, and the EC₅₀ values were calculated by GraphPad PRISM 6.

Abbreviations: GeoMean: Geometric mean; hCCR8: Human C-C motif chemokine receptor-8.

the results of omics analysis.²⁹ CCR8 is suggested to be involved in the pathogenesis of various cancer types.^{24,25} CAFs, similar to Tregs, are one of the tumor-suppressive factors known to interfere with the function of tumor immune cells by promoting fibrosis and constructing the extracellular matrix in the TME.^{1,57} CAFs with a myofibroblastic-like phenotype transfer large amounts of proteins to the surrounding endothelial cells through matrix-bound vesicles, which may contribute to cancer




Figure 5. The analysis of the binding affinity of anti-hCCR8 monoclonal antibodies for TALL-1. TALL-1 cells were suspended in 100 μ L of serially diluted C₈Mab-21 (100 μ g/mL – 0.006 μ g/mL) (A), S19017D (10 μ g/mL, 2.5 μ g/mL – 0.0006 μ g/mL) (B), or L263G8 (0.625 μ g/mL – 0.0006 μ g/mL) (C). Then, cells were treated with Alexa Fluor 488-conjugated anti-mouse immunoglobulin G. Fluorescence data were subsequently collected using the BD FACSLyric, following the calculation of the EC₅₀ by GraphPad PRISM 6.

Abbreviations: GeoMean: Geometric mean; hCCR8: Human C-C motif chemokine receptor-8.

progression and treatment resistance.^{58,59} Recently, the phenotypes of CAF have been reported to associate with either better or worse outcomes in NSCLC patients.⁶⁰ Although further functional analysis of CCR8-expressing CAFs is required, targeting CCR8 could suppress Treg and CAFs, and lead to synergistic antitumor immunotherapy results. Therefore, it is well worth evaluating the impact of C₈Mab-21 on CAFs.

5. Conclusion

C₈Mab-21, established by the CBIS method, is a useful tool for analyzing the hCCR8-positive cells by flow cytometry.

Acknowledgments

None.

Funding

This research was supported, in part, by the Japan Agency for Medical Research and Development (AMED) under Grant Nos.: JP24am0521010 (to Y.K.), JP23ama121008 (to Y.K.), JP23bm1123027 (to Y.K.), and JP23ck0106730 (to Y.K.), and by the Japan Society for the Promotion of Science (JSPS) Grants-in-Aid for Scientific Research (KAKENHI) Grant Nos.: 22K06995 (to H.S.) and 22K07224 (to Y. K.).

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

Conceptualization: Mika K. Kaneko, Yukinari Kato

Formal analysis: Tomohiro Tanaka

Investigation: Haruto Yamamoto, Yu Kaneko, Tomohiro Tanaka, Guanjie Li

Methodology: Mika K. Kaneko

Writing – original draft: Tomohiro Tanaka

Writing – review & editing: Hiroyuki Suzuki, Yukinari Kato

Ethics approval and consent to participate

All animal experiments were approved by the Animal Care and Use Committee of Tohoku University (Permit number: 2022Mda-001).

Consent for publication

Not applicable.

Availability of data

The data from this study are available in the article.

Further disclosure

The paper has been uploaded to a preprint server (DOI: 10.20944/preprints202403.1166.v1).

References

1. Bruni D, Angell HK, Galon J. The immune contexture and Immunoscore in cancer prognosis and therapeutic efficacy. *Nat Rev Cancer*. 2020;20(11):662–680.
doi: 10.1038/s41568-020-0285-7
2. Demaria O, Cornen S, Daeron M, Morel Y, Medzhitov R, Vivier E. Harnessing innate immunity in cancer therapy. *Nature*. 2019;574(7776):45–56.
doi: 10.1038/s41586-019-1593-5
3. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. *Science*. 2018;359(6382):1350–1355.
doi: 10.1126/science.aar4060
4. Karin N. Chemokines and cancer: New immune checkpoints for cancer therapy. *Curr Opin Immunol*. 2018;51:140–145.
doi: 10.1016/j.coi.2018.03.004
5. van den Bulk J, Verdegaal EM, de Miranda NF. Cancer immunotherapy: Broadening the scope of targetable tumours. *Open Biol*. 2018;8(6):180037.
doi: 10.1098/rsob.180037
6. Morad G, Helmink BA, Sharma P, Wargo JA. Hallmarks of response, resistance, and toxicity to immune checkpoint blockade. *Cell*. 2021;184(21):5309–5337.
doi: 10.1016/j.cell.2021.09.020
7. Lipson EJ, Forde PM, Hammers HJ, Emens LA, Taube JM, Topalian SL. Antagonists of PD-1 and PD-L1 in cancer treatment. *Semin Oncol*. 2015;42(4):587–600.
doi: 10.1053/j.seminoncol.2015.05.013
8. Li C, Jiang P, Wei S, Xu X, Wang J. Regulatory T cells in tumor microenvironment: New mechanisms, potential therapeutic strategies and future prospects. *Mol Cancer*. 2020;19(1):116.
doi: 10.1186/s12943-020-01234-1
9. Sharpe AH, Wherry EJ, Ahmed R, Freeman GJ. The function of programmed cell death 1 and its ligands in regulating autoimmunity and infection. *Nat Immunol*. 2007;8(3):239–245.
doi: 10.1038/ni1443
10. Yano H, Andrews LP, Workman CJ, Vignali DAA. Intratumoral regulatory T cells: Markers, subsets and their impact on anti-tumor immunity. *Immunology*. 2019;157(3):232–247.
doi: 10.1111/imm.13067
11. Ohue Y, Nishikawa H. Regulatory T (Treg) cells in cancer: Can Treg cells be a new therapeutic target? *Cancer Sci*. 2019;110(7):2080–2089.
doi: 10.1111/cas.14069
12. Tanaka A, Sakaguchi S. Regulatory T cells in cancer immunotherapy. *Cell Res*. 2017;27(1):109–118.
doi: 10.1038/cr.2016.151
13. Fridman WH, Pagès F, Sautès-Fridman C, Galon J. The immune contexture in human tumours: impact on clinical outcome. *Nat Rev Cancer*. 2012;12(4):298–306.
doi: 10.1038/nrc3245

14. Sakaguchi S. Regulatory T cells: Key controllers of immunologic self-tolerance. *Cell*. 2000;101(5):455-458.
doi: 10.1016/s0092-8674(00)80856-9
15. Brunkow ME, Jeffery EW, Hjerrild KA, *et al*. Disruption of a new forkhead/winged-helix protein, scurfy, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet*. 2001;27(1):68-73.
doi: 10.1038/83784
16. Takahashi T, Kuniyasu Y, Toda M, *et al*. Immunologic self-tolerance maintained by CD25+CD4+ naturally anergic and suppressive T cells: Induction of autoimmune disease by breaking their anergic/suppressive state. *Int Immunol*. 1998;10(12):1969-1980.
doi: 10.1093/intimm/10.12.1969
17. Tay C, Tanaka A, Sakaguchi S. Tumor-infiltrating regulatory T cells as targets of cancer immunotherapy. *Cancer Cell*. 2023;41(3):450-465.
doi: 10.1016/j.ccell.2023.02.014
18. Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol*. 2003;4(4):330-336.
doi: 10.1038/ni904
19. Mortezaee K. Selective targeting or reprogramming of intra-tumoral Tregs. *Med Oncol*. 2024;41(3):71.
doi: 10.1007/s12032-024-02300-0
20. Wakiyama H, Kato T, Furusawa A, *et al*. Treg-dominant tumor microenvironment is responsible for hyperprogressive disease after PD-1 blockade therapy. *Cancer Immunol Res*. 2022;10(11):1386-1397.
doi: 10.1158/2326-6066.Cir-22-0041
21. Guan X, Hu R, Choi Y, *et al*. Anti-TIGIT antibody improves PD-L1 blockade through myeloid and T(reg) cells. *Nature*. 2024;627:646-655.
doi: 10.1038/s41586-024-07121-9
22. Sharma P, Hu-Lieskovan S, Wargo JA, Ribas A. Primary, adaptive, and acquired resistance to cancer immunotherapy. *Cell*. 2017;168(4):707-723.
doi: 10.1016/j.cell.2017.01.017
23. Wang L, Simons DL, Lu X, *et al*. Connecting blood and intratumoral Treg cell activity in predicting future relapse in breast cancer. *Nat Immunol*. 2019;20(9):1220-1230.
doi: 10.1038/s41590-019-0429-7
24. De Simone M, Arrigoni A, Rossetti G, *et al*. Transcriptional landscape of human tissue lymphocytes unveils uniqueness of tumor-infiltrating T regulatory cells. *Immunity*. 2016;45(5):1135-1147.
doi: 10.1016/j.immuni.2016.10.021
25. Plitas G, Konopacki C, Wu K, *et al*. Regulatory T cells exhibit distinct features in human breast cancer. *Immunity*. 2016;45(5):1122-1134.
doi: 10.1016/j.immuni.2016.10.032
26. Barsheshet Y, Wildbaum G, Levy E, *et al*. CCR8(+)FOXP3(+) Treg cells as master drivers of immune regulation. *Proc Natl Acad Sci U S A*. 2017;114(23):6086-6091.
doi: 10.1073/pnas.1621280114
27. Nagira Y, Nagira M, Nagai R, *et al*. S-531011, a novel anti-human CCR8 antibody, induces potent antitumor responses through depletion of tumor-infiltrating CCR8-expressing regulatory T cells. *Mol Cancer Ther*. 2023;22(9):1063-1072.
doi: 10.1158/1535-7163.Mct-22-0570
28. Wu Y, Xi J, Li Y, *et al*. Discovery of a potent and selective CCR8 small molecular antagonist IPG7236 for the treatment of cancer. *J Med Chem*. 2023;66(7):4548-4564.
doi: 10.1021/acs.jmedchem.3c00030
29. Kim N, Kim MH, Pyo J, *et al*. CCR8 as a therapeutic novel target: Omics-integrated comprehensive analysis for systematically prioritizing indications. *Biomedicines*. 2023;11(11):2910.
doi: 10.3390/biomedicines11112910
30. Märkl F, Huynh D, Endres S, Kobold S. Utilizing chemokines in cancer immunotherapy. *Trends Cancer*. 2022;8(8):670-682.
doi: 10.1016/j.trecan.2022.04.001
31. Bernardini G, Spinetti G, Ribatti D, *et al*. I-309 binds to and activates endothelial cell functions and acts as an angiogenic molecule *in vivo*. *Blood*. 2000;96(13):4039-4045.
32. Goya I, Gutiérrez J, Varona R, *et al*. Identification of CCR8 as the specific receptor for the human beta-chemokine I-309: Cloning and molecular characterization of murine CCR8 as the receptor for TCA-3. *J Immunol*. 1998;160(4):1975-1981.
33. Liu X, Xu X, Deng W, *et al*. CCL18 enhances migration, invasion and EMT by binding CCR8 in bladder cancer cells. *Mol Med Rep*. 2019;19(3):1678-1686.
doi: 10.3892/mmr.2018.9791
34. Spinetti G, Bernardini G, Camarda G, *et al*. The chemokine receptor CCR8 mediates rescue from dexamethasone-induced apoptosis via an ERK-dependent pathway. *J Leukoc Biol*. 2003;73(1):201-207.
doi: 10.1189/jlb.0302105
35. Denis C, Deiteren K, Mortier A, *et al*. C-terminal clipping of chemokine CCL1/I-309 enhances CCR8-mediated intracellular calcium release and anti-apoptotic activity. *PLoS One*. 2012;7(3):e34199.
doi: 10.1371/journal.pone.0034199
36. Campbell JR, McDonald BR, Mesko PB, *et al*. Fc-optimized

- anti-CCR8 antibody depletes regulatory T cells in human tumor models. *Cancer Res.* 2021;81(11):2983-2994.
doi: 10.1158/0008-5472.Can-20-3585
37. Saito M, Harigae Y, Li G, *et al.* C(3)Mab-2: An anti-mouse CCR3 monoclonal antibody for immunocytochemistry. *Monoclon Antib Immunodiagn Immunother.* 2022;41(1):45-49.
doi: 10.1089/mab.2021.0050
 38. Saito M, Suzuki H, Tanaka T, *et al.* Development of an anti-mouse CCR8 monoclonal antibody (C(8)Mab-1) for flow cytometry and immunocytochemistry. *Monoclon Antib Immunodiagn Immunother.* 2022;41(6):333-338.
doi: 10.1089/mab.2021.0069
 39. Nanamiya R, Takei J, Asano T, *et al.* Development of anti-human CC chemokine receptor 9 monoclonal antibodies for flow cytometry. *Monoclon Antib Immunodiagn Immunother.* 2021;40(3):101-106.
doi: 10.1089/mab.2021.0007
 40. Ouchida T, Suzuki H, Tanaka T, Kaneko MK, Kato Y. Cx(4)Mab-1: A novel anti-mouse CXCR4 monoclonal antibody for flow cytometry. *Monoclon Antib Immunodiagn Immunother.* 2024;43(1):10-16.
doi: 10.1089/mab.2023.0023
 41. Schöneberg T, Liebscher I. Mutations in G protein-coupled receptors: Mechanisms, pathophysiology and potential therapeutic approaches. *Pharmacol Rev.* 2021;73(1):89-119.
doi: 10.1124/pharmrev.120.000011
 42. Hauser AS, Attwood MM, Rask-Andersen M, Schiöth HB, Gloriam DE. Trends in GPCR drug discovery: New agents, targets and indications. *Nat Rev Drug Discov.* 2017;16(12):829-842.
doi: 10.1038/nrd.2017.178
 43. Proudfoot AE. Chemokine receptors: Multifaceted therapeutic targets. *Nat Rev Immunol.* 2002;2(2):106-115.
doi: 10.1038/nri722
 44. Jo M, Jung ST. Engineering therapeutic antibodies targeting G-protein-coupled receptors. *Exp Mol Med.* 2016;48(2):e207.
doi: 10.1038/emm.2015.105
 45. Itai S, Yamada S, Kaneko MK, *et al.* Establishment of EMab-134, a sensitive and specific anti-epidermal growth factor receptor monoclonal antibody for detecting squamous cell carcinoma cells of the oral cavity. *Monoclon Antib Immunodiagn Immunother.* 2017;36(6):272-281.
doi: 10.1089/mab.2017.0042
 46. Asano T, Ohishi T, Takei J, *et al.* AntiHER3 monoclonal antibody exerts antitumor activity in a mouse model of colorectal adenocarcinoma. *Oncol Rep.* 2021;46(2):173.
doi: 10.3892/or.2021.8124
 47. Tanaka T, Ohishi T, Asano T, *et al.* An antiTROP2 monoclonal antibody TrMab6 exerts antitumor activity in breast cancer mouse xenograft models. *Oncol Rep.* 2021;46(1):132.
doi: 10.3892/or.2021.8083
 48. Kudo Y, Suzuki H, Tanaka T, Kaneko MK, Kato Y. Development of a novel anti-CD44 variant 5 monoclonal antibody C(44)Mab-3 for multiple applications against pancreatic carcinomas. *Antibodies (Basel).* 2023;12(2):31.
doi: 10.3390/antib12020031
 49. Suzuki H, Kaneko MK, Kato Y. Roles of podoplanin in malignant progression of tumor. *Cells.* 2022;11(3):575.
doi: 10.3390/cells11030575
 50. Kaneko MK, Suzuki H, Kato Y. Establishment of a novel cancer-specific anti-HER2 monoclonal antibody H(2)Mab-250/H(2)CasMab-2 for breast cancers. *Monoclon Antib Immunodiagn Immunother.* 2024;43(2):35-43.
doi: 10.1089/mab.2023.0033
 51. Arimori T, Mihara E, Suzuki H, *et al.* Locally misfolded HER2 expressed on cancer cells is a promising target for development of cancer-specific antibodies. *Structure.* 2024;32:536-549.e5.
doi: 10.1016/j.str.2024.02.007
 52. Ueyama A, Nogami W, Nashiki K, *et al.* Immunotherapy targeting CCR8+ regulatory T cells induces antitumor effects via dramatic changes to the intratumor CD8+ T cell profile. *J Immunol.* 2023;211(4):673-682.
doi: 10.4049/jimmunol.2300067
 53. Kidani Y, Nogami W, Yasumizu Y, *et al.* CCR8-targeted specific depletion of clonally expanded Treg cells in tumor tissues evokes potent tumor immunity with long-lasting memory. *Proc Natl Acad Sci USA.* 2022;119(7):e2114282119.
doi: 10.1073/pnas.2114282119
 54. Zheng D, Wang X, Cheng L, *et al.* The chemokine receptor CCR8 is a target of chimeric antigen T cells for treating T cell malignancies. *Front Immunol.* 2022;13:808347.
doi: 10.3389/fimmu.2022.808347
 55. Li G, Suzuki H, Ohishi T, *et al.* Antitumor activities of a defucosylated antiEpCAM monoclonal antibody in colorectal carcinoma xenograft models. *Int J Mol Med.* 2023;51(2):18.
doi: 10.3892/ijmm.2023.5221
 56. Niwa R, Satoh M. The current status and prospects of antibody engineering for therapeutic use: focus on glycoengineering technology. *J Pharm Sci.* 2015;104(3):930-941.
doi: 10.1002/jps.24316
 57. Kalluri R. The biology and function of fibroblasts in cancer. *Nat Rev Cancer.* 2016;16(9):582-598.

- doi: 10.1038/nrc.2016.73
58. Santi A, Kay EJ, Neilson LJ, *et al.* Cancer-associated fibroblasts produce matrix-bound vesicles that influence endothelial cell function. *Sci Signal.* 2024;17(827):eade0580.
doi: 10.1126/scisignal.ade0580
 59. Eskandari-Malayeri F, Rezaei M. Immune checkpoint inhibitors as mediators for immunosuppression by cancer-associated fibroblasts: A comprehensive review. *Front Immunol.* 2022;13:996145.
doi: 10.3389/fimmu.2022.996145
 60. Cords L, Engler S, Haberecker M, *et al.* Cancer-associated fibroblast phenotypes are associated with patient outcome in non-small cell lung cancer. *Cancer Cell.* 2024;42(3):396-412.e395.
doi: 10.1016/j.ccell.2023.12.021