

Communication

C₈Mab-21: A Novel Anti-Human CCR8 Monoclonal Antibody for Flow Cytometry

Tomohiro Tanaka, Hiroyuki Suzuki, Guanjie Li, Mika K. Kaneko, Yukinari Kato *

Department of Antibody Drug Development, Tohoku University Graduate School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8575, Japan

* Correspondence: yukinari.kato.e6@tohoku.ac.jp (Y.K.); Tel.: +81-22-717-8207

Abstract: C-C motif chemokine receptor-8 (CCR8) belongs to class A of G protein-coupled receptors (GPCRs). CCR8 interacts with the specific chemokine ligand CCL1/I-309 in humans, which is produced by various cells, including tumor-associated macrophages and regulatory T cells (Treg). CCR8 is highly expressed on Treg and T-helper 2 (T_H2) cells recruited to the inflammation site and is implicated in allergy, asthma, and cancer progression. The CCR8⁺Treg has been suggested to be an important regulator in the immunosuppressive tumor microenvironment (TME); therefore, it has been desired to develop sensitive monoclonal antibodies (mAbs) for CCR8. This study developed a specific mAb for human CCR8 (hCCR8), which is useful for flow cytometry by employing the Cell-Based Immunization and Screening (CBIS) method. The established anti-hCCR8 mAb, C₈Mab-21, (mouse IgM, kappa) reacted with hCCR8-overexpressed Chinese hamster ovary-K1 (CHO/hCCR8) cells, TALL-1 (acute T lymphoblastic leukemia), CCRF-HSB2 (human T-lymphoblastic leukemia), and natural killer cells, which express endogenous hCCR8 by flow cytometry. Furthermore, C₈Mab-21 demonstrated a moderate binding affinity for CHO/hCCR8 and TALL-1 with a dissociation constant of 6.5×10^{-8} M and 2.0×10^{-8} M, respectively. C₈Mab-21, which was established by the CBIS method, could be a useful tool for analyzing the hCCR8-related biological response using flow cytometry.

Keywords: CCR8; CBIS method; monoclonal antibody; flow cytometry

1. Introduction

Immune checkpoint inhibitors have become effective and powerful strategies for cancer therapy.[1] In particular, the development of antibody drugs targeting immune checkpoint molecules, such as programmed-cell death-1 (PD-1), cytotoxic T lymphocyte antigen 4 (CTLA-4), and PD-1 ligand 1 (PD-L1), has achieved remarkable therapeutic results.[2-4] PD-1 inhibits the excessive activation of conventional T cells by suppressing costimulatory signaling and renders them dysfunctional or exhausted.[5] PD-1 and CTLA-4 are also expressed in regulatory T cells (Treg), one of the immunosuppressors in the tumor microenvironment (TME). The inhibition could potentiate the activation and immunosuppressive function of Treg cells.[6]

Treg is defined as a CD4⁺T cell with CD25 and Foxp3 and maintains self-tolerance to prevent excessive immune responses and autoimmune diseases.[7] Treg suppresses the effector functions of T cells through the secretion of immunosuppressive cytokines, such as interleukin-10, transforming growth factor- β , and cytotoxic granzyme/perforin.[7-11] Intratumoral Treg suppresses antitumor T cell responses and thus resists the effects of immune checkpoint inhibitor therapy.[12,13] Antibodies against T cell immunoreceptors with Ig and ITIM domains (TIGIT), one of the immune checkpoint molecules, improve the effectiveness of PD-L1 antibodies by suppressing Treg.[14] Therefore, the development of immunotherapy targeting Treg is expected.[6,15]

Intratumoral Treg expresses a high level of C-C motif chemokine receptor-8 (CCR8). Additionally, the CCR8-expressing Treg has also increased the expression of CD25 and Foxp3 compared with CCR8-negative Treg.[16] Therefore, CCR8-expressing Treg is considered to have potent immunosuppressive functions. The CCR8-expressing Treg is known to be correlated with poor prognosis in some cancer patients.[17,18] CCR8 is attractive as a target molecule for the next

cancer immunotherapy.[19] Anti-CCR8 drugs, including S-531011,[20] IPG7236,[21] and SRF114[22] are undergoing clinical trials.

CCR8 is one of the seven transmembrane-spanning G protein-coupled receptors (GPCRs). Human CCR8 is known to bind to five C-C chemokine ligands (CCLs): CCL1/I-309, CCL4, CCL16, CCL17, and CCL18.[23] CCR8 is upregulated in not only Treg, but also various cancers, including breast, non-small cell lung (NSCLC), bladder, and colorectal cancer.[17,18] CCR8 mediates bladder cancer cell migration, invasion, and epithelial-mesenchymal transition by interacting with CCL18.[24] CCR8 and its specific ligand CCL1/I-309 regulate the immune system, which mediates the progression of diseases such as cancers. CCR8-CCL1/I-309 axis promotes migration and inhibits apoptosis in Treg and lymphomas.[25,26] Therefore, CCR8-targeting antibodies will contribute to the elucidation of pathological mechanisms, diagnosis, and therapy.

Using the Cell-Based Immunization and Screening (CBIS) method, we previously developed numerous monoclonal antibodies (mAbs) against chemokine receptors, including mouse CCR3,[27,28] mouse CCR8,[29-31] human CCR9,[32] and mouse CXCR4.[33] In this study, we have successfully developed an anti-human CCR8 (hCCR8) mAb using the CBIS method, which is applicable to flow cytometry.

2. Materials and Methods

2.1. Preparation of Cell Lines

LN229, Chinese hamster ovary (CHO)-K1, and P3X63Ag8U.1 (P3U1) cells were obtained from the American Type Culture Collection (Manassas, VA). TALL-1 and CCRF-HSB2 cells were obtained from the Japanese Collection of Research Bioresources (JCRB) Cell Bank (Osaka, Japan). The human natural killer (NK) cells (donor lot. 4022602, purity > 70%) were purchased from Takara Bio (Shiga, Japan). pCMV6neo-myc-DDK vector with hCCR8 (Accession No.: NM_005201) was purchased from OriGene Technologies, Inc. (Rockville, MD). The plasmid was transfected into the cell lines using a Neon transfection system (Thermo Fisher Scientific, Inc., Waltham, MA). Subsequently, LN229 and CHO-K1, which stably overexpressed hCCR8 with C-terminal myc-DDK tags (hereinafter described as LN229/hCCR8 and CHO/hCCR8, respectively) were established using a cell sorter (SH800; Sony Corp., Tokyo, Japan), following cultivation in a medium containing 0.5 mg/mL G418 (Nacalai Tesque, Inc., Kyoto, Japan).

2.2. Production of Hybridomas

For developing anti-hCCR8 mAbs, two female 6-week-old BALB/c mice were immunized intraperitoneally with 1×10^8 cells of LN229/hCCR8. The immunogen was harvested after brief exposure to 1 mM ethylenediaminetetraacetic acid (EDTA; Nacalai Tesque, Inc.). We added Imject Alum Adjuvant (Thermo Fisher Scientific Inc.) as an adjuvant in the first immunization. Three additional injections of 1×10^8 cells of LN229/hCCR8 were performed without an adjuvant every week. We performed a final booster immunization of 1×10^8 cells of LN229/hCCR8 intraperitoneally two days before harvesting splenocytes from mice. We conducted cell-fusion of the harvested splenocytes with P3U1 cells using polyethylene glycol 1500 (PEG1500; Roche Diagnostics, Indianapolis, IN).

2.3. Flow Cytometric Analyses

CHO-K1 and CHO/hCCR8 cells were harvested after brief exposure to 1 mM EDTA. CHO-K1, CHO/hCCR8, TALL-1, and CCRF-HSB2 cells were washed with 0.1% bovine serum albumin in phosphate-buffered saline and treated with primary mAbs for 30 min at 4°C. Afterward, cells were treated with Alexa Fluor 488-conjugated anti-mouse IgG (1:1000) following the collection of fluorescence data, using the SA3800 Cell Analyzer (Sony Corp.). The anti-human CD198 (CCR8) mAbs (clones S19017D and L263G8) were purchased from BioLegend (San Diego, CA). The Alexa Fluor 488-conjugated anti-mouse IgG was purchased from Cell Signaling Technology, Inc. (Danvers, MA).

2.4. Determination of the Binding Affinity by Flow Cytometry

CHO/hCCR8 and TALL-1 were suspended in 100 μ L serially diluted of CsMab-21 (100 μ g/mL to 0.006 μ g/mL), S19017D (10 μ g/mL to 0.0006 μ g/mL for CHO/hCCR8, 10 μ g/mL, 2.5 μ g/mL to 0.0006 μ g/mL for TALL-1), or L263G8 (10 μ g/mL to 0.0006 μ g/mL for CHO/hCCR8, 0.625 μ g/mL to 0.0006 μ g/mL for TALL-1), after which Alexa Fluor 488-conjugated anti-mouse IgG (1:200) was added. Fluorescence data were subsequently collected, using the BD FACSLyric (BD Biosciences, Franklin Lakes, NJ), following the calculation of the dissociation constant (K_D) by fitting the binding isotherms into the built-in; one-site binding model in GraphPad PRISM 6 (GraphPad Software, Inc., La Jolla, CA).

3. Results

3.1. Establishment of Anti-hCCR8 mAbs by the CBIS Method

To develop anti-hCCR8 mAbs, we employed the CBIS method using hCCR8-overexpressed cells. Anti-hCCR8 mAbs-producing hybridoma screening was conducted using flow cytometry (Figure 1). Two mice were intraperitoneally immunized with LN229/hCCR8 weekly for a total of 5 times. Subsequently, hybridomas were seeded into 96-well plates, after which flow cytometric analysis was used to select CHO/hCCR8-reactive and CHO-K1-nonreactive supernatants of hybridomas. Afterward, we obtained CHO/hCCR8-reactive supernatant in only 1 of 956 wells (0.10%). We finally established clone CsMab-21 (mouse IgM, kappa) by limiting dilution and additional screening.




Figure 1. A schematic procedure of anti-hCCR8 mAbs production. The procedure of the Cell-Based Immunization and Screening (CBIS) method for antibody development. (A) LN229/hCCR8 cells were immunized into two mice using intraperitoneal injection. (B) The spleen cells from mice were fused with P3U1 myeloma cells. (C) The culture supernatants of hybridoma were screened by flow cytometry using CHO-K1 and CHO/hCCR8. (D) After limiting dilution of hybridomas and additional analysis, CsMab-21 was finally established.

3.2. Flow Cytometric Analysis

Flow cytometric analysis was conducted using C₈Mab-21 and commercially available anti-human CD198 (CCR8) mAbs (clone S19017D and L263G8) against CHO-K1, CHO/hCCR8, TALL-1, CCRF-HSB2, and NK cells. Results showed that C₈Mab-21, S19017D, and L263G8 recognize CHO/hCCR8 dose-dependently (Figure 2A). C₈Mab-21, and L263G8 did not react with parental CHO-K1 cells even at 20 $\mu\text{g}/\text{mL}$ of mAbs. S19017D slightly reacted to CHO-K1 cells at 20 $\mu\text{g}/\text{mL}$ and also 2 $\mu\text{g}/\text{mL}$ (Figure 2B). Regarding endogenously hCCR8-expressing cells in Figure 3, C₈Mab-21 recognized TALL-1, CCRF-HSB2, and NK cells at a concentration of 2 $\mu\text{g}/\text{mL}$ or 20 $\mu\text{g}/\text{mL}$ (Figure 3). S19017D reacted to TALL-1 and CCRF-HSB2 dose-dependently even at a concentration of 0.02 $\mu\text{g}/\text{mL}$. However, NK cells were not recognized by S19017D even at a concentration of 20 $\mu\text{g}/\text{mL}$ (Figure 3). L263G8 reacted to TALL-1 and CCRF-HSB2 even at 0.02 $\mu\text{g}/\text{mL}$, and also NK cells at a concentration of 2 $\mu\text{g}/\text{mL}$ or higher of mAb (Figure 3). C₈Mab-21 could detect exogenously and endogenously expressed naïve conformational hCCR8 in flow cytometry.




Figure 2. Flow cytometric analysis of anti-hCCR8 mAbs against CHO/hCCR8 and CHO-K1. CHO/hCCR8 (A) and CHO-K1 cells (B) were treated with 0.02–20 $\mu\text{g}/\text{mL}$ of C₈Mab-21, S19017D, and L263G8 (red line), followed by treatment with Alexa Fluor 488-conjugated anti-mouse IgG. Fluorescence data were collected using the SA3800 Cell Analyzer. Black line, control (no primary antibody treatment).




Figure 3. Flow cytometric analysis of anti-hCCR8 mAbs against endogenously hCCR8-expressing cells. TALL-1 (A), CCRF-HSB2 (B), and NK cells (C) were treated with 0.02–20 µg/mL of CsMab-21, S19017D, and L263G8 (red line), followed by treatment with Alexa Fluor 488-conjugated anti-mouse IgG. Fluorescence data were collected using the SA3800 Cell Analyzer. Black line, control (no primary antibody treatment).

3.3. Determination of the Binding Affinity of Anti-hCCR8 mAbs to CHO/hCCR8

The binding affinity of CsMab-21, S19017D, and L263G8 was assessed with exogenously hCCR8-expressed CHO/hCCR8 using flow cytometry. Results showed that the K_D values of CsMab-21, S19017D, and L263G8 for CHO/hCCR8 are 6.5×10^{-8} M, 2.6×10^{-9} M, and 1.2×10^{-9} M, respectively (Figure 4). These results indicate that CsMab-21 possesses a moderate affinity for exogenously overexpressed hCCR8 in CHO-K1 cells.




Figure 4. The analysis of the binding affinity of anti-hCCR8 mAbs for CHO/hCCR8. CHO/hCCR8 cells were suspended in 100 μL of serially diluted C8Mab-21 (100 $\mu\text{g}/\text{mL}$ to 0.006 $\mu\text{g}/\text{mL}$) (A), S19017D (10 $\mu\text{g}/\text{mL}$ to 0.0006 $\mu\text{g}/\text{mL}$) (B), or L263G8 (10 $\mu\text{g}/\text{mL}$ to 0.0006 $\mu\text{g}/\text{mL}$) (C). Then, cells were treated with Alexa Fluor 488-conjugated anti-mouse IgG. Fluorescence data were subsequently collected using the BD FACSLyric, following the calculation of the dissociation constant (K_D) by GraphPad PRISM 6.

3.4. Determination of the Binding Affinity of Anti-hCCR8 mAbs to TALL-1

The binding affinity of C8Mab-21, S19017D, and L263G8 was analyzed with endogenously hCCR8-expressing TALL-1 using flow cytometry. Results indicated that the K_D values of C8Mab-21, S19017D, and L263G8 for TALL-1 are $2.0 \times 10^{-8} \text{ M}$, $4.6 \times 10^{-10} \text{ M}$, and $7.8 \times 10^{-11} \text{ M}$, respectively (Figure 5). These results showed that C8Mab-21 possesses a moderate affinity for endogenously expressing hCCR8 in TALL-1 leukemia cells.




Figure 5. The analysis of the binding affinity of anti-hCCR8 mAbs for TALL-1. TALL-1 cells were suspended in 100 μL of serially diluted C8Mab-21 (100 $\mu\text{g}/\text{mL}$ to 0.006 $\mu\text{g}/\text{mL}$) (A), S19017D (10 $\mu\text{g}/\text{mL}$, 2.5 $\mu\text{g}/\text{mL}$ to 0.0006 $\mu\text{g}/\text{mL}$) (B), or L263G8 (0.625 $\mu\text{g}/\text{mL}$ to 0.0006 $\mu\text{g}/\text{mL}$) (C). Then, cells were treated with Alexa Fluor 488-conjugated anti-mouse IgG. Fluorescence data were subsequently collected using the BD FACSLyric, following the calculation of the dissociation constant (K_D) by GraphPad PRISM 6.

4. Discussion

GPCRs, including chemokine receptors, are focused as targets for many diseases, including inflammatory disorders and cancers.[34,35] GPCRs transmit signals to intracellular molecules about extracellular conditions and govern broad cell dynamics, such as proliferation, homeostasis, migration, and motility of the cells.[34,35] Therapeutic drugs, including mAbs targeting GPCRs, have been developed to date; however, the complexity of the structure, the small area of epitope regions, and the difficulty of purifying the protein as an immunogen pose high hurdles.[36,37]

Unlike protein purifications, the preparation of antigens is not complicated in the CBIS method. Furthermore, it is possible to retain the antigen structure and modification such as glycosylation and folding using the CBIS method. We have successfully developed multiple mAbs by the CBIS method against human epidermal growth factor receptor 1 (HER1; EGFR),[38] HER3,[39] trophoblast cell surface antigen 2,[40] epithelial cell adhesion molecule,[41] CD44,[42-44] and podoplanin.[45] Furthermore, some of the developed mAbs by the CBIS method exhibited cancer specificity by recognizing unique cancer-specific epitopes.[46-51] Therefore, the CBIS method is an efficient and useful tactic for generating diverse antibodies targeting membrane proteins.

In the immunosuppressive TME, CD8⁺T cells are exhausted along with the induction of CCR8⁺Treg. CCR8⁺Treg infiltration is confirmed to be associated with high TOX, an exhaustion marker of T cells, expression in CD8⁺T cells in some types of cancer patients.[52,53] Elimination of CCR8⁺Treg using antibodies is expected to advance the treatment of these cancer patients. Targeting CCR8 might be more specific in antitumor activity than other approaches aimed at Treg removal.[53,54] In mice, CCR8⁺T cell depletion therapy using anti-CCR8 mAbs induces tumor-specific immune responses without triggering autoimmune responses or immune reactions in TME.[55] Since C₈Mab-21 could recognize cell surface hCCR8, we plan to investigate the function of C₈Mab-21 against Treg, such as detection and interfering effects in future studies. Furthermore, we have previously enhanced antibody-dependent cellular cytotoxicity (ADCC) activity and complement-dependent cytotoxicity (CDC) by modifying isotypes and defucosylation in mAbs.[56,57] Since C₈Mab-21 is mouse IgM, which has no ADCC activity, it will be converted into a mouse IgG_{2a} version to evaluate the effect of antitumor activities in xenograft models.

Interestingly, a correlation between cancer-associated fibroblasts (CAFs) and CCR8 has been found from the results of omics analysis.[22] CCR8 is suggested to be involved in the pathogenesis of various cancer types. CAFs are one of the tumor suppressor factors, the same as Treg, that are known to interfere with the function of tumor immune cells by promoting fibrosis and constructing an extracellular matrix in the TME.[58,59] CAFs with a myofibroblastic-like phenotype transfer large amounts of proteins to the surrounding endothelial cells through matrix-bound vesicles, which might contribute to cancer progression and treatment resistance.[60,61] Recently, the phenotypes of CAF have been reported to be associated with better or worse outcomes in NSCLC patients.[62] Although further functional analysis of CCR8-expressing CAFs is required, targeting CCR8 might suppress Treg and CAFs, and lead to synergistic antitumor immunotherapy results. In that case, it is well worth evaluating the impact of C₈Mab-21 on CAFs as well in future studies.

Authorship confirmation/contribution statement: T.T. and G.L. performed the experiments. M.K.K. and Y.K. designed the experiments. T.T. analyzed the data. T.T., H.S., and Y.K. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Author Disclosure Statement: The authors have no conflict of interest.

Funding Information: This research was supported, in part, by Japan Agency for Medical Research and Development (AMED) under Grant Nos.: JP23ama121008 (to Y.K.), JP23am0401013 (to Y.K.), JP23bm1123027 (to Y.K.), and JP23ck0106730 (to Y.K.), and by the Japan Society for the Promotion of Science (JSPS) Grants-in-Aid for Scientific Research (KAKENHI) Grant Nos.: 21K20789 (to T.T.), 22K06995 (to H.S.), 21K07168 (to M.K.K.), and 22K07224 (to Y.K.).

Institutional Review Board Statement: The animal study protocol was approved by the Animal Care and Use Committee of Tohoku University (Permit number: 2022MdA-001) for studies involving animals.

References

1. Ribas, A.; Wolchok, J.D. Cancer immunotherapy using checkpoint blockade. *Science* **2018**, *359*, 1350-1355, doi:10.1126/science.aar4060.
2. van den Bulk, J.; Verdegaal, E.M.; de Miranda, N.F. Cancer immunotherapy: broadening the scope of targetable tumours. *Open Biol* **2018**, *8*, doi:10.1098/rsob.180037.
3. Morad, G.; Helmink, B.A.; Sharma, P.; Wargo, J.A. Hallmarks of response, resistance, and toxicity to immune checkpoint blockade. *Cell* **2021**, *184*, 5309-5337, doi:10.1016/j.cell.2021.09.020.

4. Lipson, E.J.; Forde, P.M.; Hammers, H.J.; Emens, L.A.; Taube, J.M.; Topalian, S.L. Antagonists of PD-1 and PD-L1 in Cancer Treatment. *Semin Oncol* **2015**, *42*, 587-600, doi:10.1053/j.seminoncol.2015.05.013.
5. Li, C.; Jiang, P.; Wei, S.; Xu, X.; Wang, J. Regulatory T cells in tumor microenvironment: new mechanisms, potential therapeutic strategies and future prospects. *Mol Cancer* **2020**, *19*, 116, doi:10.1186/s12943-020-01234-1.
6. Ohue, Y.; Nishikawa, H. Regulatory T (Treg) cells in cancer: Can Treg cells be a new therapeutic target? *Cancer Sci* **2019**, *110*, 2080-2089, doi:10.1111/cas.14069.
7. Sakaguchi, S. Regulatory T cells: key controllers of immunologic self-tolerance. *Cell* **2000**, *101*, 455-458, doi:10.1016/s0092-8674(00)80856-9.
8. Brunkow, M.E.; Jeffery, E.W.; Hjerrild, K.A.; Paepfer, B.; Clark, L.B.; Yasayko, S.A.; Wilkinson, J.E.; Galas, D.; Ziegler, S.F.; Ramsdell, F. Disruption of a new forkhead/winged-helix protein, scurfy, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet* **2001**, *27*, 68-73, doi:10.1038/83784.
9. Takahashi, T.; Kuniyasu, Y.; Toda, M.; Sakaguchi, N.; Itoh, M.; Iwata, M.; Shimizu, J.; Sakaguchi, S. Immunologic self-tolerance maintained by CD25+CD4+ naturally anergic and suppressive T cells: induction of autoimmune disease by breaking their anergic/suppressive state. *Int Immunol* **1998**, *10*, 1969-1980, doi:10.1093/intimm/10.12.1969.
10. Tay, C.; Tanaka, A.; Sakaguchi, S. Tumor-infiltrating regulatory T cells as targets of cancer immunotherapy. *Cancer Cell* **2023**, *41*, 450-465, doi:10.1016/j.ccell.2023.02.014.
11. Fontenot, J.D.; Gavin, M.A.; Rudensky, A.Y. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol* **2003**, *4*, 330-336, doi:10.1038/ni904.
12. Mortezaee, K. Selective targeting or reprogramming of intra-tumoral Tregs. *Med Oncol* **2024**, *41*, 71, doi:10.1007/s12032-024-02300-0.
13. Wakiyama, H.; Kato, T.; Furusawa, A.; Okada, R.; Inagaki, F.; Furumoto, H.; Fukushima, H.; Okuyama, S.; Choyke, P.L.; Kobayashi, H. Treg-Dominant Tumor Microenvironment Is Responsible for Hyperprogressive Disease after PD-1 Blockade Therapy. *Cancer Immunol Res* **2022**, *10*, 1386-1397, doi:10.1158/2326-6066.Cir-22-0041.
14. Guan, X.; Hu, R.; Choi, Y.; Srivats, S.; Nabet, B.Y.; Silva, J.; McGinnis, L.; Hendricks, R.; Nutsch, K.; Banta, K.L.; et al. Anti-TIGIT antibody improves PD-L1 blockade through myeloid and T(reg) cells. *Nature* **2024**, doi:10.1038/s41586-024-07121-9.
15. Sharma, P.; Hu-Lieskovan, S.; Wargo, J.A.; Ribas, A. Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell* **2017**, *168*, 707-723, doi:10.1016/j.cell.2017.01.017.
16. Wang, L.; Simons, D.L.; Lu, X.; Tu, T.Y.; Solomon, S.; Wang, R.; Rosario, A.; Avalos, C.; Schmolze, D.; Yim, J.; et al. Connecting blood and intratumoral Treg cell activity in predicting future relapse in breast cancer. *Nat Immunol* **2019**, *20*, 1220-1230, doi:10.1038/s41590-019-0429-7.
17. De Simone, M.; Arrigoni, A.; Rossetti, G.; Gruarin, P.; Ranzani, V.; Politano, C.; Bonnal, R.J.P.; Provasi, E.; Sarnicola, M.L.; Panzeri, I.; et al. Transcriptional Landscape of Human Tissue Lymphocytes Unveils Uniqueness of Tumor-Infiltrating T Regulatory Cells. *Immunity* **2016**, *45*, 1135-1147, doi:10.1016/j.immuni.2016.10.021.
18. Plitas, G.; Konopacki, C.; Wu, K.; Bos, P.D.; Morrow, M.; Putintseva, E.V.; Chudakov, D.M.; Rudensky, A.Y. Regulatory T Cells Exhibit Distinct Features in Human Breast Cancer. *Immunity* **2016**, *45*, 1122-1134, doi:10.1016/j.immuni.2016.10.032.
19. Barsheshet, Y.; Wildbaum, G.; Levy, E.; Vitenshtein, A.; Akinseye, C.; Griggs, J.; Lira, S.A.; Karin, N. CCR8(+)/FOXP3(+) Treg cells as master drivers of immune regulation. *Proc Natl Acad Sci U S A* **2017**, *114*, 6086-6091, doi:10.1073/pnas.1621280114.
20. Nagira, Y.; Nagira, M.; Nagai, R.; Nogami, W.; Hirata, M.; Ueyama, A.; Yoshida, T.; Yoshikawa, M.; Shinonome, S.; Yoshida, H.; et al. S-531011, a Novel Anti-Human CCR8 Antibody, Induces Potent Antitumor Responses through Depletion of Tumor-Infiltrating CCR8-Expressing Regulatory T Cells. *Mol Cancer Ther* **2023**, *22*, 1063-1072, doi:10.1158/1535-7163.Mct-22-0570.
21. Wu, Y.; Xi, J.; Li, Y.; Li, Z.; Zhang, Y.; Wang, J.; Fan, G.H. Discovery of a Potent and Selective CCR8 Small Molecular Antagonist IPG7236 for the Treatment of Cancer. *J Med Chem* **2023**, *66*, 4548-4564, doi:10.1021/acs.jmedchem.3c00030.
22. Kim, N.; Kim, M.H.; Pyo, J.; Lee, S.M.; Jang, J.S.; Lee, D.W.; Kim, K.W. CCR8 as a Therapeutic Novel Target: Omics-Integrated Comprehensive Analysis for Systematically Prioritizing Indications. *Biomedicines* **2023**, *11*, doi:10.3390/biomedicines11112910.
23. Märkl, F.; Huynh, D.; Endres, S.; Kobold, S. Utilizing chemokines in cancer immunotherapy. *Trends Cancer* **2022**, *8*, 670-682, doi:10.1016/j.trecan.2022.04.001.
24. Liu, X.; Xu, X.; Deng, W.; Huang, M.; Wu, Y.; Zhou, Z.; Zhu, K.; Wang, Y.; Cheng, X.; Zhou, X.; et al. CCL18 enhances migration, invasion and EMT by binding CCR8 in bladder cancer cells. *Mol Med Rep* **2019**, *19*, 1678-1686, doi:10.3892/mmr.2018.9791.

25. Spinetti, G.; Bernardini, G.; Camarda, G.; Mangoni, A.; Santoni, A.; Capogrossi, M.C.; Napolitano, M. The chemokine receptor CCR8 mediates rescue from dexamethasone-induced apoptosis via an ERK-dependent pathway. *J Leukoc Biol* **2003**, *73*, 201-207, doi:10.1189/jlb.0302105.
26. Denis, C.; Deiteren, K.; Mortier, A.; Tounsi, A.; Fransen, E.; Proost, P.; Renauld, J.C.; Lambeir, A.M. C-terminal clipping of chemokine CCL1/I-309 enhances CCR8-mediated intracellular calcium release and anti-apoptotic activity. *PLoS One* **2012**, *7*, e34199, doi:10.1371/journal.pone.0034199.
27. Saito, M.; Harigae, Y.; Li, G.; Asano, T.; Tanaka, T.; Suzuki, H.; Kaneko, M.K.; Kato, Y. C(3)Mab-2: An Anti-Mouse CCR3 Monoclonal Antibody for Immunocytochemistry. *Monoclon Antib Immunodiagn Immunother* **2022**, *41*, 45-49, doi:10.1089/mab.2021.0050.
28. Asano, T.; Suzuki, H.; Tanaka, T.; Saito, M.; Li, G.; Goto, N.; Nanamiya, R.; Kaneko, M.K.; Kato, Y. C(3)Mab-3: A Monoclonal Antibody for Mouse CC Chemokine Receptor 3 for Flow Cytometry. *Monoclon Antib Immunodiagn Immunother* **2022**, *41*, 74-79, doi:10.1089/mab.2021.0062.
29. Saito, M.; Suzuki, H.; Tanaka, T.; Asano, T.; Kaneko, M.K.; Kato, Y. Development of an Anti-Mouse CCR8 Monoclonal Antibody (C(8)Mab-1) for Flow Cytometry and Immunocytochemistry. *Monoclon Antib Immunodiagn Immunother* **2022**, *41*, 333-338, doi:10.1089/mab.2021.0069.
30. Tanaka, T.; Nanamiya, R.; Takei, J.; Nakamura, T.; Yanaka, M.; Hosono, H.; Sano, M.; Asano, T.; Kaneko, M.K.; Kato, Y. Development of Anti-Mouse CC Chemokine Receptor 8 Monoclonal Antibodies for Flow Cytometry. *Monoclon Antib Immunodiagn Immunother* **2021**, *40*, 65-70, doi:10.1089/mab.2021.0005.
31. Suzuki, H.; Saito, M.; Asano, T.; Tanaka, T.; Kitamura, K.; Kudo, Y.; Kaneko, M.K.; Kato, Y. C(8)Mab-3: An Anti-Mouse CCR8 Monoclonal Antibody for Immunocytochemistry. *Monoclon Antib Immunodiagn Immunother* **2022**, *41*, 110-114, doi:10.1089/mab.2022.0002.
32. Nanamiya, R.; Takei, J.; Asano, T.; Tanaka, T.; Sano, M.; Nakamura, T.; Yanaka, M.; Hosono, H.; Kaneko, M.K.; Kato, Y. Development of Anti-Human CC Chemokine Receptor 9 Monoclonal Antibodies for Flow Cytometry. *Monoclon Antib Immunodiagn Immunother* **2021**, *40*, 101-106, doi:10.1089/mab.2021.0007.
33. Ouchida, T.; Suzuki, H.; Tanaka, T.; Kaneko, M.K.; Kato, Y. Cx(4)Mab-1: A Novel Anti-Mouse CXCR4 Monoclonal Antibody for Flow Cytometry. *Monoclon Antib Immunodiagn Immunother* **2024**, *43*, 10-16, doi:10.1089/mab.2023.0023.
34. Proudfoot, A.E. Chemokine receptors: multifaceted therapeutic targets. *Nat Rev Immunol* **2002**, *2*, 106-115, doi:10.1038/nri722.
35. Schöneberg, T.; Liebscher, I. Mutations in G Protein-Coupled Receptors: Mechanisms, Pathophysiology and Potential Therapeutic Approaches. *Pharmacol Rev* **2021**, *73*, 89-119, doi:10.1124/pharmrev.120.000011.
36. Jo, M.; Jung, S.T. Engineering therapeutic antibodies targeting G-protein-coupled receptors. *Exp Mol Med* **2016**, *48*, e207, doi:10.1038/emm.2015.105.
37. Hauser, A.S.; Attwood, M.M.; Rask-Andersen, M.; Schiöth, H.B.; Gloriam, D.E. Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug Discov* **2017**, *16*, 829-842, doi:10.1038/nrd.2017.178.
38. Itai, S.; Yamada, S.; Kaneko, M.K.; Chang, Y.W.; Harada, H.; Kato, Y. Establishment of EMab-134, a Sensitive and Specific Anti-Epidermal Growth Factor Receptor Monoclonal Antibody for Detecting Squamous Cell Carcinoma Cells of the Oral Cavity. *Monoclon Antib Immunodiagn Immunother* **2017**, *36*, 272-281, doi:10.1089/mab.2017.0042.
39. Asano, T.; Ohishi, T.; Takei, J.; Nakamura, T.; Nanamiya, R.; Hosono, H.; Tanaka, T.; Sano, M.; Harada, H.; Kawada, M.; et al. Anti-HER3 monoclonal antibody exerts antitumor activity in a mouse model of colorectal adenocarcinoma. *Oncol Rep* **2021**, *46*, doi:10.3892/or.2021.8124.
40. Tanaka, T.; Ohishi, T.; Asano, T.; Takei, J.; Nanamiya, R.; Hosono, H.; Sano, M.; Harada, H.; Kawada, M.; Kaneko, M.K.; et al. An anti-TROP2 monoclonal antibody TrMab-6 exerts antitumor activity in breast cancer mouse xenograft models. *Oncol Rep* **2021**, *46*, doi:10.3892/or.2021.8083.
41. Kaneko, M.K.; Ohishi, T.; Takei, J.; Sano, M.; Nakamura, T.; Hosono, H.; Yanaka, M.; Asano, T.; Sayama, Y.; Harada, H.; et al. Anti-EpCAM monoclonal antibody exerts antitumor activity against oral squamous cell carcinomas. *Oncol Rep* **2020**, *44*, 2517-2526, doi:10.3892/or.2020.7808.
42. Kudo, Y.; Suzuki, H.; Tanaka, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 5 Monoclonal Antibody C(44)Mab-3 for Multiple Applications against Pancreatic Carcinomas. *Antibodies (Basel)* **2023**, *12*, doi:10.3390/antib12020031.
43. Suzuki, H.; Ozawa, K.; Tanaka, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 7/8 Monoclonal Antibody, C(44)Mab-34, for Multiple Applications against Oral Carcinomas. *Biomedicines* **2023**, *11*, doi:10.3390/biomedicines11041099.
44. Ejima, R.; Suzuki, H.; Tanaka, T.; Asano, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 6 Monoclonal Antibody C(44)Mab-9 for Multiple Applications against Colorectal Carcinomas. *Int J Mol Sci* **2023**, *24*, doi:10.3390/ijms24044007.
45. Suzuki, H.; Kaneko, M.K.; Kato, Y. Roles of Podoplanin in Malignant Progression of Tumor. *Cells* **2022**, *11*, doi:10.3390/cells11030575.

46. Kaneko, M.K.; Suzuki, H.; Ohishi, T.; Nakamura, T.; Tanaka, T.; Kato, Y. A Cancer-Specific Monoclonal Antibody against HER2 Exerts Antitumor Activities in Human Breast Cancer Xenograft Models. *Int J Mol Sci* **2024**, *25*, doi:10.3390/ijms25031941.
47. Kaneko, M.K.; Suzuki, H.; Kato, Y. Establishment of a Novel Cancer-specific Anti-HER2 Monoclonal Antibody H2Mab-250/H2CasMab-2 for breast cancers. *Monoclon Antib Immunodiagn Immunother* **2024**, in press.
48. Arimori, T.; Mihara, E.; Suzuki, H.; Ohishi, T.; Tanaka, T.; Kaneko, M.K.; Takagi, J.; Kato, Y. Locally misfolded HER2 expressed on cancer cells is a promising target for development of cancer-specific antibodies. *Structure* **2024**, doi:10.1016/j.str.2024.02.007.
49. Suzuki, H.; Ohishi, T.; Tanaka, T.; Kaneko, M.K.; Kato, Y. A Cancer-Specific Monoclonal Antibody against Podocalyxin Exerted Antitumor Activities in Pancreatic Cancer Xenografts. *Int J Mol Sci* **2023**, *25*, doi:10.3390/ijms25010161.
50. Suzuki, H.; Ohishi, T.; Kaneko, M.K.; Kato, Y. A Humanized and Defucosylated Antibody against Podoplanin (humLpMab-23-f) Exerts Antitumor Activities in Human Lung Cancer and Glioblastoma Xenograft Models. *Cancers (Basel)* **2023**, *15*, doi:10.3390/cancers15205080.
51. Kato, Y.; Kaneko, M.K. A cancer-specific monoclonal antibody recognizes the aberrantly glycosylated podoplanin. *Sci Rep* **2014**, *4*, 5924, doi:10.1038/srep05924.
52. Ueyama, A.; Nogami, W.; Nashiki, K.; Haruna, M.; Miwa, H.; Hagiwara, M.; Nagira, M.; Wada, H.; Nagira, Y. Immunotherapy Targeting CCR8+ Regulatory T Cells Induces Antitumor Effects via Dramatic Changes to the Intratumor CD8+ T Cell Profile. *J Immunol* **2023**, *211*, 673-682, doi:10.4049/jimmunol.2300067.
53. Kidani, Y.; Nogami, W.; Yasumizu, Y.; Kawashima, A.; Tanaka, A.; Sonoda, Y.; Tona, Y.; Nashiki, K.; Matsumoto, R.; Hagiwara, M.; et al. CCR8-targeted specific depletion of clonally expanded Treg cells in tumor tissues evokes potent tumor immunity with long-lasting memory. *Proc Natl Acad Sci U S A* **2022**, *119*, doi:10.1073/pnas.2114282119.
54. Zheng, D.; Wang, X.; Cheng, L.; Qin, L.; Jiang, Z.; Zhao, R.; Li, Y.; Shi, J.; Wu, Q.; Long, Y.; et al. The Chemokine Receptor CCR8 Is a Target of Chimeric Antigen T Cells for Treating T Cell Malignancies. *Front Immunol* **2022**, *13*, 808347, doi:10.3389/fimmu.2022.808347.
55. Campbell, J.R.; McDonald, B.R.; Mesko, P.B.; Siemers, N.O.; Singh, P.B.; Selby, M.; Sproul, T.W.; Korman, A.J.; Vlach, L.M.; Houser, J.; et al. Fc-Optimized Anti-CCR8 Antibody Depletes Regulatory T Cells in Human Tumor Models. *Cancer Res* **2021**, *81*, 2983-2994, doi:10.1158/0008-5472.Can-20-3585.
56. Li, G.; Suzuki, H.; Ohishi, T.; Asano, T.; Tanaka, T.; Yanaka, M.; Nakamura, T.; Yoshikawa, T.; Kawada, M.; Kaneko, M.K.; et al. Antitumor activities of a defucosylated anti-EpCAM monoclonal antibody in colorectal carcinoma xenograft models. *Int J Mol Med* **2023**, *51*, doi:10.3892/ijmm.2023.5221.
57. Tanaka, T.; Suzuki, H.; Ohishi, T.; Kaneko, M.K.; Kato, Y. Antitumor activities against breast cancers by an afucosylated anti-HER2 monoclonal antibody H(2) Mab-77-mG(2a) -f. *Cancer Sci* **2024**, *115*, 298-309, doi:10.1111/cas.16008.
58. Bruni, D.; Angell, H.K.; Galon, J. The immune contexture and Immunoscore in cancer prognosis and therapeutic efficacy. *Nat Rev Cancer* **2020**, *20*, 662-680, doi:10.1038/s41568-020-0285-7.
59. Kalluri, R. The biology and function of fibroblasts in cancer. *Nat Rev Cancer* **2016**, *16*, 582-598, doi:10.1038/nrc.2016.73.
60. Santi, A.; Kay, E.J.; Neilson, L.J.; McGarry, L.; Lilla, S.; Mullin, M.; Paul, N.R.; Fercoq, F.; Koulouras, G.; Rodriguez Blanco, G.; et al. Cancer-associated fibroblasts produce matrix-bound vesicles that influence endothelial cell function. *Sci Signal* **2024**, *17*, eade0580, doi:10.1126/scisignal.ade0580.
61. Eskandari-Malayeri, F.; Rezaei, M. Immune checkpoint inhibitors as mediators for immunosuppression by cancer-associated fibroblasts: A comprehensive review. *Front Immunol* **2022**, *13*, 996145, doi:10.3389/fimmu.2022.996145.
62. Cords, L.; Engler, S.; Haberecker, M.; Rüschoff, J.H.; Moch, H.; de Souza, N.; Bodenmiller, B. Cancer-associated fibroblast phenotypes are associated with patient outcome in non-small cell lung cancer. *Cancer Cell* **2024**, *42*, 396-412.e395, doi:10.1016/j.ccell.2023.12.021.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.